


475-129

4-8-1976

1/1

Best Available Copy

REDUCED
40%**SNAROL**
PELLETSPUNCH
HERE**SNAROL**
PELLETS**KILLS**
SLUGS & SNAILS**SNAROL**
PELLETS**SNAROL**
PELLETS**KILLS**
SLUGS & SNAILS

SNAROL PELLETS are a specially prepared bait for the control of snails and slugs. The Metolachlor in SNAROL PELLETS attracts and kills snails and slugs. These pests are most active in and around damp cool places, such as flower and vegetable gardens, around fences, hedgerows and trees. The presence of snails and slugs can be determined by their shiny mucus-like trails.

DIRECTIONS

Spread SNAROL PELLETS late in the day before or after thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL PELLETS to apply them to the soil and not directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, cress, tomatoes, apply to soil surface, around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous wetting and high temperatures cause SNAROL PELLETS gradually to lose some of their effectiveness, so during spring and summer months, repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baits, ten days apart, usually gives control of new broods. During late and winter treatment every 2 weeks is usually satisfactory.



WARNING KEEP OUT OF REACH OF CHILDREN.
THIS PESTICIDE MAY BE FATAL TO CHILDREN, AND
DOGS OR OTHER PETS IF EATEN. KEEP CHILDREN
AND PETS OUT OF TREATED AREA.

ACTIVE INGREDIENT: Metolachlor 1.2%
INERT INGREDIENTS 98.8%

NET WT. 2-1/2 LBS.**SPREAD BEFORE
OR AFTER SPRINKLING**

One pound of SNAROL PELLETS is sufficient to treat about 1200 sq ft of surface area. SNAROL is also available in PEARL form.

WARNING

May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Harmful dust. Avoid breathing dust, contact with skin, eyes and clothing. Wash hands with soap and water after using.

Keep out of reach of children. Keep dogs and other pets, domestic animals or poultry away from treated areas.

Do not apply directly on fruits, vegetables or crops, but apply on ground under plants. Dispose of empty container or any spillage promptly—burn or bury. Store in a cool, dry place. This product is toxic to wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BOYLE-MIDWAY INC.

NEW YORK, NEW YORK 10017


EPA Registration No. 475-129-AI

EPA Est. 475-CA-1

© Boyle Midway Co. Made in U.S.A. SP 13

ACCEPTED

APR 8 1976

Under the
F.F.D.
CH. 100
FOGELSON
EPA Reg. No. 475-129**SNAROL**
PELLETSG
H
I
J
K
L
M
N
O
P
Q
R
S
T
U
V
W
X
Y
Z