

✓
LABEL AMENDMENT APPLICATION OF JUNE 17, 1987 FOR
EPA REG. 475-129 TO MODIFY INGREDIENT STATEMENT
(LOWER LEVEL OF ACTIVE INGREDIENT -- FORMULA AMEND-
MENT FILED MARCH 31, 1987). NO CHANGES ARE PROPOSED
OTHER THAN IN THE INGREDIENT STATEMENT FROM MOST RE-
CENT EPA APPROVED LABEL COPY (JULY 2, 1985).

FRONT PANEL

From the makers of Black Flag

SNAROL
SNAIL & SLUG
KILLER PELLETS

ACCEPTED
AUG 26 1987
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended for the pesticide
registered under
EPA Reg. No. 475-129

Spread on soil surface around fruits and vegetables,
flowers and ornamental plants.
This box sufficiently treats about 3,000 square feet.

Active Ingredients: Metaldehyde.....2.5%
Inert Ingredients.....97.5%
Total.....100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING

THIS PESTICIDE MAY BE FATAL TO CHILDREN AND DOGS
OR OTHER PETS IF EATEN. KEEP CHILDREN AND PETS
OUT OF TREATED AREA.
See side panel for additional precautionary
statements.

Net Wt. 2-1/2 lb. (1.13 kg)

NOTE: For 5 lb. box substitute the following where appropriate:
"6,000" square feet
"5 lb. (2.26 kg)"

Snarol & Slug Killer Pellets

RIGHT SIDE PANEL

**SNAROL
SNAIL & SLUG
KILLER PELLETS**

Spread before or after sprinkling

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed or harmful if inhaled. Avoid breathing dust, contact with skin, eyes and clothing. Wash hands with soap and water after using. Keep dogs and other pets, domestic animals or poultry away from treated areas.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Do not apply directly on fruits, vegetables or crops but apply on ground under plants. This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Boyle-Midway, Inc.
New York, New York 10017

EPA Registration No. 475-129
EPA Est. 475-CA-1

Snarol Snail & Slug Killer Pellets

LEFT SIDE PANEL

**PUNCH
HERE**

**SNAROL
SNAIL & SLUG
KILLER PELLETS**

**SPREAD BEFORE OR
AFTER SPRINKLING**

One pound of SNAROL PELLETS is sufficient to treat about 1,200 sq. ft. of surface area. SNAROL is also available in MEAL form.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

Spread SNAROL PELLETS late in the day, before or after thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL PELLETS to apply it to the soil area under plants, do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.

STORAGE AND DISPOSAL

Store in a cool, dry place away from children. Do not reuse empty container. Wrap container and put in trash collection.

Snarol Snail & Slug Killer Pellets

REAR PANEL

HOW CAN I BE CERTAIN I HAVE SNAILS OR SLUGS?

You'll probably never see them directly because these pests feed at night. But just look for the obvious clues such as holes in tomatoes, damaged or dying flowers, or shiny mucous-like trails in your garden or flower bed, under boards or flower pots, around fences, hedges and around bushes.

DOES SNAROL WORK?

The Metaldehyde in Snarol attracts and kills snails and slugs.

WHERE DO I USE SNAROL?

Late in the day before or after watering infested areas. For strawberries, don't use within 6 days of harvesting.

HOW OFTEN SHOULD I USE SNAROL?

During spring and summer, repeat treatment every week or ten days (three of such baitings will usually kill the offspring, too). During fall and winter, treatment every two weeks is usually satisfactory.

