

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

NOV 2 1989

Mr. John Pazzola
Boyle-Hidway Home-Ford Products, Inc.
South Avenue & Hale Street
Cransford, NJ 07016

Dear Mr. Pazzola:

Subject: Response to Metaldehyde Registration Standard
Snarol Snail & Slug Killer Pellets
EPA Registration No. 475-129
Your Submissions Dated July 11 and September 14, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes listed below before you release the product for shipment bearing the amended label.

1. Revise the Precautionary Statement to read:

Avoid all spray, dust, or dust breathing, dust.
Avoid contact with skin, eyes, or clothing.
Wash hands with soap. Wash contaminated
clothing before reuse. Keep kids and other
pets out of treated areas.

2. Revise the Statement of Practical Treatment. Use the following format:

If Swallowed: _____
If Inhaled: _____
If On Skin: _____
If In Eyes: _____

STWOB:11/24/89:UC-7:KALCO:11/21/89:12/12/89:CI:VO:CT

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

EPA Form 1320-1 (12-70) OFFICIAL FILE COPY

-2-

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

This will also acknowledge receipt of the Generic Data Exemption Statement. This information will be placed in the file for this product.

Sincerely yours,

Dennis H. Edwards, Jr.
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

Re and
to 75%

Response to "Re-registration of
Pesticide Products Containing
Metaldehyde as the Active Ingredient"


EPA Reg. No 475-129

Submitted by Boyle-Midway Household
Products, Inc. 9/14/89

Changes encircled in red

SNAROL

SNAIL & SLUG KILLER PELLETS




From the makers of



SNAROL

SNAIL & SLUG KILLER PELLETS



SPREAD BEFORE OR
AFTER SPRINKLING

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS

WARNING

May be fatal if swallowed or harm-
ful if inhaled. Avoid breathing dust
contact with skin, eyes and cloth-
ing. Wash hands with soap and
water after using. Keep dogs and
other pets, domestic animals or
poultry away from treated areas.

STATEMENT OF PRACTICAL TREATMENT
If swallowed. Call a physician or
Poison Control Center. Drink 1 or
2 glasses of water and induce
vomiting by touching back of
throat with finger. Do not induce
vomiting or give anything by
mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Do not apply directly on fruits, veg-
etables, or crops but apply on ground
under plants. Do not apply
directly to water or wetlands (swamps,
bays, marshes and pebbles). Do not
contaminate water when disposing of
equipment washwater. Apply this pro-
duct only as specified on this label.

STORAGE AND DISPOSAL

Store in a cool, dry place away
from children. Do not reuse empty
container. Wrap container and put
in trash collection.

Distributed by

BOYLE-MIDWAY
HOUSEHOLD PRODUCTS, INC.
NEW YORK, N.Y. 10017
EPA Registration No. 475-129
EPA Est. 567-CA1
PO Box 100000 New York, NY 10010

ACCEPTED
with COMMENTS
in EPA Letter 8806

IV 2 1989

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registration No. 475-129.

SNAROL

SNAIL & SLUG KILLER PELLETS



SAVE
30%
ON ANY
BLACK FLAG
PRODUCT
COUPON INSIDE

SPREAD ON SOIL SURFACE AROUND FRUITS AND
VEGETABLES, FLOWERS AND ORNAMENTAL PLANTS.

This box sufficiently treats about 2,700 square feet.

ACTIVE INGREDIENTS: Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-tetraoxycyclo-octane).....	2.5%
INERT INGREDIENTS.....	97.5%
TOTAL.....	100.0%

KEEP OUT OF REACH OF CHILDREN
WARNING

THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN
KEEP PETS OUT OF TREATED AREA.

See side panel for additional precautionary statements.

NET WT 2 1/4 LB (1.02 kg)

Front & Back Panels: Black, White, Yellow, Red type on White, Green Background

Side Panels:
Black, Yellow, Green type
on White, Green Background

SNAROL

SNAIL & SLUG KILLER PELLETS



PUNCH
HERE

SNAROL[®]

SNAIL & SLUG KILLER PELLETS



SPREAD BEFORE OR
AFTER SPRINKLING

One pound of SNAROL PELLETS is sufficient to treat about 1,200 sq. ft. of surface area. SNAROL is also available in MEAL form.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Spread SNAROL PELLETS late in the day, after it is thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL PELLETS to apply it to the soil area under plants—do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.

SNAROL[®]


SNAIL & SLUG KILLER PELLETS

HOW CAN I BE CERTAIN I HAVE SNAILS OR SLUGS?

You'll probably never see them directly because these pests feed at night. But just look for the obvious clues such as holes in tomatoes, damaged or dying flowers, or shiny mucous-like trails in your garden or flower bed, under boards or flower pots, around fences, hedges and around bushes.


HOW DOES SNAROL WORK?

The Metaldehyde in Snarol attracts and kills snails and slugs.




WHERE DO I USE SNAROL?

Spread Snarol on the soil surface, not directly on the plants or edible parts. Snarol can be used around fruit and vegetable plants.



WHEN DO I USE SNAROL?

Late in the day after watering infested areas. For strawberries, don't use within 6 days of harvesting.



HOW OFTEN SHOULD I USE SNAROL?

During spring and summer, repeat treatment every week or 12-14 days (three of such baitings will usually kill the offspring, too). During fall and winter, treatment every two weeks is usually satisfactory.