

475-130

9-19-1973

1/2

ACCEPTED

SEP 19 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 475-130

**PUNCH
HERE**

SNAROL MEAL

KILLS SNAILS & SLUGS

SNAROL is a specially prepared bait for the control of snails and slugs. The metaldehyde in SNAROL attracts and kills snails and slugs. These pests are night feeders and inhabit damp, cool places such as flower and vegetable gardens, around fences, hedges and bushes. The presence of snails and slugs can be determined by their shiny mucous-like trails.

DIRECTIONS

Apply SNAROL late in the day, after thoroughly wetting the areas likely to be infested. Spread SNAROL lightly over infested area. Care should be taken when spreading SNAROL to apply it to the soil area under plants, do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.

2001

2/2

SNAROL MEAL



SNAROL MEAL



KILLS SNAILS & SLUGS

WARNING: KEEP OUT OF REACH OF CHILDREN
Read warning and directions on side panel before using.

ACTIVE INGREDIENT: Metaldehyde 3.0%
INERT INGREDIENTS 97.0%

NET WT. 2-1/2 LBS.

SNAROL MEAL



SNAROL MEAL KILLS SNAILS & SLUGS

SPREAD BEFORE OR
AFTER SPRINKLING

One pound of SNAROL MEAL is sufficient to treat about 1200 sq. ft. of surface area. SNAROL is also available in PELLET form.

WARNING

May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Harmful dust. Avoid contact with skin, eyes and clothing. Wash hands with soap and water after using.

Keep out of reach of children.

Keep domestic animals and poultry off treated area. Bait is attractive to dogs and chickens.

Do not apply directly on fruits, vegetables or crops, but apply on ground under plants. Dispose of empty container or any spillage promptly, burn or bury. Store in a cool, dry place.

BOYLE-MIDWAY INC.
NEW YORK, N.Y. 10017
© Boyle-Midway Div.

EPA Registration No. 475-130-AA
Made in U.S.A. SM-7