

475-130 11-21-1974

1/2

# SNAROL MEAL



# SNAROL<sup>®</sup> MEAL



# KILLS SNAILS & SLUGS

**WARNING** KEEP OUT OF REACH OF CHILDREN.  
THIS PESTICIDE MAY BE FATAL TO CHILDREN, AND  
DOGS OR OTHER PETS IF EATEN. KEEP CHILDREN  
AND PETS OUT OF TREATED AREA.  
Read and follow all directions. Read warning statement on side panel.

ACTIVE INGREDIENT: Metaldehyde ..... 3.0%  
INERT INGREDIENTS ..... 97.0%

**NET WT. 2-1/2 LBS.**



## SNAROL MEAL KILLS SNAILS & SLUGS

**SPREAD BEFORE OR  
AFTER SPRINKLING**

One pound of SNAROL MEAL is sufficient to treat about 1200 sq. ft. of surface area. SNAROL is also available in PELLET form.

May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Harmful dust. Avoid breathing dust, contact with skin, eyes and clothing. Wash hands with soap and water after using.

Keep out of reach of children. Keep domestic animals and poultry off treated area. Bait is attractive to dogs and chickens. Do not apply directly on fruits, vegetables or crops, but apply on ground under plants. Do not re-use empty container. Dispose of empty container or any spillage promptly—burn or bury. Store in a cool, dry place.

This product is toxic to wildlife and birds. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

BOYLE-MIDWAY INC.  
NEW YORK, NEW YORK 10017  
EPA Registration No. 475-130-AA  
EPA Est. 475-CA-1  
Boyle-Midway Div. Made in U.S.A. SM-8

# SNAROL MEAL



F ■  
G ■  
H ■  
I ■  
J ■  
K ■

ACCEPTED  
495-130  
NOV 21 1974

2/2

**PUNCH  
HERE**

# **SNAROL**

## **MEAL**

# **KILLS SNAILS & SLUGS**

SNAROL is a specially prepared bait for the control of snails and slugs. The metaldehyde in SNAROL attracts and kills snails and slugs. These pests are night feeders and inhabit damp, cool places such as flower and vegetable gardens, around fences, hedges and bushes. The presence of snails and slugs can be determined by their shiny mucous-like trails.

### **DIRECTIONS**

Apply SNAROL late in the day, after thoroughly wetting the areas likely to be infested. Spread SNAROL lightly over infested area. Care should be taken when spreading SNAROL to apply it to the soil area under plants, do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.