

475-130

5-31-1979


1/2

# SNAROL MEAL



# SNAROL

**MEAL** From the makers of 



# SNAROL MEAL

**KILLS  
SNAILS & SLUGS**

**KILLS  
SNAILS & SLUGS**

**CAN BE USED AROUND  
FRUITS &  
VEGETABLES**

### WARNING

May be fatal if swallowed. If swallowed, induce vomiting. Call physician immediately. Harmful dust. Avoid breathing dust, contact with skin, eyes and clothing. Wash hands with soap and water after using.

Keep out of reach of children. Keep domestic animals and poultry off treated area. Bait is attractive to dogs and chickens. Do not apply directly on fruits, vegetables or crops, but apply on ground under plants. Do not reuse empty container. Dispose of empty container or any spillage promptly—burn or bury. Store in a cool, dry place. This product is toxic to birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BOYLE-MIDWAY INC  
NEW YORK, NEW YORK 10017  
EPA Registration No. 475-130-04  
EPA Est. 475-344  
© Boyle-Midway Div. Made in U.S.A. SM-12-Z

# SNAROL MEAL



069421-00004  
MAY 31 1979  
Under the Federal Insecticide, Fungicide, and Rodenticide Act  
475-130  
7/8/79


# "YOU MAY HAVE THIS PROBLEM"



Holes in Tomatoes



Damaged & Dying Flowers



## SNAROL MEAL KILLS SNAILS & SLUGS

# SNAROL kills snails and slugs

### HOW CAN I BE CERTAIN I HAVE SNAILS OR SLUGS?

You'll probably never see them directly because these pests feed at night. But just look for the clues above. Another clue is shiny mucous-like trails in your garden or flowerbed, under boards or flower pots, around fences, hedges and around bushes.

### HOW DOES SNAROL WORK?



The Metaldehyde in Snarol attracts and kills snails and slugs.


### WHERE DO I USE SNAROL?

Spread Snarol on the soil surface, not directly on the plants or edible parts. Snarol can be used around fruit and vegetable plants.



### WHEN DO I USE SNAROL?

Late in the day after watering infested areas. For strawberries, don't use within 6 days of harvesting.



### HOW OFTEN SHOULD I USE SNAROL?

During spring and summer, repeat treatment every week or ten days; three of such baitings will usually kill the offspring, too! During fall and winter, treatment every two weeks is usually satisfactory.

One pound of SNAROL MEAL is sufficient to treat about 1200 sq. ft. of surface area. SNAROL is also available in PELLET form.

### DIRECTIONS

Spread SNAROL MEAL late in the day, after thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL MEAL to apply it to the soil area under plants, do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.