

✓
LABEL AMENDMENT APPLICATION OF JUNE 17, 1987 FOR EPA REG. 475-130
TO MODIFY INGREDIENT STATEMENT (LOWER LEVEL OF ACTIVE -- FORMULA
AMENDMENT FILED MARCH 31, 1987). NO CHANGES ARE PROPOSED OTHER
THAN IN THE INGREDIENT STATEMENT FROM MOST RECENT EPA APPROVED
LABEL COPY (July 26, 1985)

FRONT PANEL

From the makers of Black Flag

SNAROL
SNAIL & SLUG
KILLER MEAL

Spread on soil surface around fruits and vegetables, flowers and
ornamental plants.
This box sufficiently treats about 3,000 square feet.


Active Ingredient: Metaldehyde.....2.5%
Inert Ingredients.....97.5%
Total.....100.0%

ACCEPTED
AUG 26 1987
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 475-130

KEEP OUT OF REACH OF CHILDREN


WARNING

THIS PESTICIDE MAY BE FATAL TO CHILDREN AND DOGS OR OTHER PETS
IF EATEN. KEEP CHILDREN AND PETS OUT OF TREATED AREA
See side panel for additional precautionary statements



Net Wt. 2-1/2 lb. (1.13 kg)

NOTE: For 5 lb. box substitute the following where appropriate:
"6,000" square feet
"5 lb. (2.26 kg)"



44-183
p. 2 of 4

Proposed Label Copy
for
Snarol Snail & Slug Killer Meal

RIGHT SIDE PANEL

SNAROL
SNAIL & SLUG
KILLER MEAL

Spread after sprinkling

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed or harmful if inhaled. Avoid breathing dust, contact with skin, eyes and clothing. Wash hands with soap and water after using. Keep dogs and other pets, domestic animals or poultry away from treated areas.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Do not apply directly on fruits, vegetables or crops but apply on ground under plants. This product is toxic to wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Boyle-Midway, Inc.
New York, New York 10017

EPA Registration No. 475-130
EPA Est. 475-CA-1

Proposed Label Copy

For

Snarol & Snail & Slug Killer Meal

LEFT SIDE PANEL

PUNCH
HERE

SNAROL
SNAIL & SLUG
KILLER MEAL

USE AFTER SPRINKLING

One pound of SNAROL MEAL is sufficient to treat about 1,200 sq. ft. of surface area. SNAROL is also available in PELLET form.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Spread SNAROL MEAL late in the day, after thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL to apply it to the soil area under plants, do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. Three baitings ten days apart, usually gives control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.

STORAGE AND DISPOSAL

Store in a cool, dry place away from children. Do not reuse empty container. Wrap container and put in trash collection.

Proposed Label Copy
for
Snarol Snail & Slug Killer Meal

REAR PANEL

HOW CAN I BE CERTAIN I HAVE SNAILS OR SLUGS?

You'll probably never see them directly because these pests feed at night. But just look for the obvious clues such as holes in tomatoes, damaged or dying flowers, or shiny mucous-like trails in your garden or flower bed, under boards or flower pots, around fences, hedges and around bushes.

DOES SNAROL WORK?

The Metaldehyde in Snarol attracts and kills snails and slugs.

WHERE DO I USE SNAROL?

Late in the day after watering infested areas. For strawberries, don't use within 6 days of harvesting.

HOW OFTEN SHOULD I USE SNAROL?

During spring and summer, repeat treatment every week or ten days (three of such baitings will usually kill the offspring, too). During fall and winter, treatment every two weeks is usually satisfactory.

