


Revised Label Copy
 "Snarol Meal-Kills Snails & Slugs"
 EPA Reg. No. 475-130
 Submitted by Boyle-Midway House-
 hold Products, Inc.
 3/19/90
 Changes noted in red.

SNAIL & SLUG KILLER MEAL

From the makers of

**BLACK
FLAG**

SNAROL SNAIL & SLUG KILLER MEAL




SPREAD AFTER SPRINKLING

ACCEPTED
 JUN 1 1990
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 475-130

SNAROL[®]

SNAIL & SLUG KILLER MEAL



SPREAD ON SOIL SURFACE AROUND FRUITS AND
 VEGETABLES, FLOWERS AND ORNAMENTAL PLANTS.

This box sufficiently treats about 2,700 square feet.

ACTIVE INGREDIENTS: Metaldehyde (2,4,6,8-tetramethyl-1,3,5,7-
 tetraoxycyclo-octane) 2.9%
 INERT INGREDIENTS 97.5%
 TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

WARNING

THIS PESTICIDE MAY BE FATAL TO DOGS OR OTHER PETS IF EATEN.

KEEP PETS OUT OF TREATED AREA.

See side panel for additional precautionary statements.

NET WT 2 1/4 LB (1.02 Kg)

**SAVE
30¢**

ON ANY
BLACK FLAG
PRODUCT
COUPON INSID

SNAROL SNAIL & SLUG KILLER MEAL



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Harmful if swallowed. Avoid
 breathing dust. Avoid contact
 with skin, eyes, or clothing.
 Wash hands with soap. Wash
 contaminated clothing before
 reuse. Keep dogs and other
 pets out of treated areas.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician or Poison
 Control Center. Drink 1 or
 2 glasses of water and induce
 vomiting by touching back of
 throat with finger. Do not
 induce vomiting or give any-
 thing by mouth to an un-
 conscious person.
 If On Skin: Wash with soap and water.
 If In Eyes: Flush with plenty of water.
 Get medical attention if
 irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly on fruits, veg-
 etables or crops but apply on ground
 under plants. Do not apply
 directly to water or wetlands (swamps,
 bays, marshes and potholes). Do not
 contaminate water when disposing of
 equipment washwaters. Apply this pro-
 duct only as specified on this label.

STORAGE AND DISPOSAL

Store in a cool, dry place away from chil-
 dren. Do not reuse empty container. Wrap
 container and put in trash collection.

Boyle-Midway
 HOUSEHOLD PRODUCTS, INC.
 NEW YORK, N.Y. 10017

(EPA Registration No. 475-130)

(N) (S) (C) (A)

USE OF THIS PRODUCT IS UNLAWFUL IN CANADA

PUNCH
HERE

SNAROL[®] SNAIL & SLUG KILLER MEAL



SPREAD AFTER SPRINKLING

One pound of SNAROL MEAL is sufficient to treat about 1,200 sq. ft. of surface area. SNAROL is also available in PELLET form.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Spread SNAROL MEAL late in the day, after thoroughly wetting the areas likely to be infested. Care should be taken when spreading SNAROL MEAL to apply it to the soil area under plants—do not apply directly on plants.

For cabbage, melons, carrots, celery, kale, peas, beans, corn, tomatoes, apply to soil surface around plants. Do not contaminate edible parts. For strawberries, do not use within 6 days of harvesting.

Continuous watering and high temperatures cause SNAROL gradually to lose some of its effectiveness, so during spring and summer months repeat treatment every week or ten days. Such repeated treatment will also serve to control new broods emerging during this period. These baitings ten days apart, usually give control of new hatch. During fall and winter, treatment every 2 weeks is usually satisfactory.


SNAROL[®] SNAIL & SLUG KILLER MEAL

HOW CAN I BE CERTAIN I HAVE SNAILS OR SLUGS?

You'll probably never see them directly because these pests feed at night. But just look for obvious clues such as holes in tomatoes, damaged or dying flowers, or shiny mucous-like trails in your garden or flower bed, under boards or flower pots, around fences, hedges and around bushes.

HOW DOES SNAROL WORK?

The Metaldehyde in Snarol attracts and kills snails and slugs.




WHEN DO I USE SNAROL?

Late in the day after watering infested areas. For strawberries, don't use within 6 days of harvesting.



WHERE DO I USE SNAROL?

Spread Snarol on the soil surface, not directly on the plants or edible parts. Snarol can be used around fruit and vegetable plants.



HOW OFTEN SHOULD I USE SNAROL?

During spring and summer, repeat treatment every week or ten days (three of such baitings will usually kill the offspring, too). During fall and winter, treatment every two weeks is usually satisfactory.

