



# TARTOX<sup>®</sup>

**AGRICULTURAL INSECTICIDE  
WETTABLE POWDER**



**POISON**



**ACTIVE INGREDIENT:**

Antimonyl Potassium Tartrate (Tartar Emetic).....99%  
INERT INGREDIENTS..... 1%  
Antimony expressed as metallic....36.1%

**DIRECTIONS FOR USE**

Measure the specified amount of this product, then wash through filler screen in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low-volume, gear-type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proved physically compatible and safe to plants. Emulsifiable concentrates should always be thoroughly emulsified with at least half of total water before adding this product.

**RECOMMENDATIONS**—(ALSO READ USE PRECAUTIONS & DIRECTIONS FOR USE)

Maximum permissible rate per acre, expressed as Tartox, is given in parenthesis (), after each crop claim.

**CITRUS** (Grapefruit, Lemons, Oranges, Tangerines)--For control of ~~citrus thrips~~ apply 1/2 lb. TARTOX with 1/2 lb. of sugar in 20-50 gal. water per acre by a blower type spray rig. If a conventional or boom rig is used, apply 2 lb. each TARTOX and sugar in 200 gal. water per acre.

For control of snails mix 2 lb. TARTOX with 4 lb. sugar in 100 gal. water and apply 3-4 gal. spray per average tree. Do not apply within 30 days of harvest. (2 1/2 lb.)

**GLADIOLUS**--For control of thrips mix 2 lb. of TARTOX with 4 lb. of sugar in 100 gal. water and apply when thrips first appear. For best control apply first spray before the flower spikes form and repeat at weekly intervals as needed during the growing season.

100 LB. NET

**DANGER**

**Keep Out of Reach of Children**

**MAY BE FATAL IF SWALLOWED.**

Do not breathe dust or spray mist. ✓  
Do not get in eyes, on skin or on clothing.  
Wash thoroughly after handling.  
Do not contaminate food or feed.

Call a physician IMMEDIATELY. Until he arrives follow the directions given here.

Contains ANTIMONY (Tartar Emetic)

ANTIDOTE--If conscious, induce vomiting by giving a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Give milk or white of eggs beaten with water. Keep patient warm and quiet.

**OBSERVE FOLLOWING USE PRECAUTIONS**

Observe use limitations given on the label for specific crops.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damage caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing, number and rate of applications needed will vary with local conditions.

Toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Apply only to areas specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-37

S-SF-700317-WC

**STAUFFER CHEMICAL COMPANY** NEW YORK, NY 10017

S-SF 681101

**BEFORE USING STUDY SAFETY DIRECTIONS—DESTROY EMPTY CONTAINERS—BURY OR BURN—STAY OUT OF SMOKE AND FUMES**  
**BEFORE USING STUDY SAFETY DIRECTIONS—DESTROY EMPTY CONTAINERS—BURY OR BURN—STAY OUT OF SMOKE AND FUMES**