


COPPER SULFATE MONOHYDRATE 20 DUST

AN AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENT:

COPPER SULFATE MONOHYDRATE 20%

(7.16% copper expressed as metallic)

INERT INGREDIENTS 80%

100%

DIRECTION FOR USE

Except as otherwise noted, apply 20-40 lbs. per acre when the air is calm using ground power equipment or aircraft. For best results, apply before evidence of disease or as specified on the label for specific crops. Unless otherwise shown, repeat applications at 7-10 day intervals to maintain a light film of dust on foliage during periods of infection.

Before marketing food crops, remove residues by washing, stripping, wiping, or other effective means.

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED.
AVOID INHALING DUST.
AVOID CONTACT WITH SKIN.

OBSERVE FOLLOWING USE PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fibre or pasture crops. **The grower is responsible** for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service, as the timing, number, and rate of applications needed will vary with local conditions.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Made in U.S.A.
Stauffer Chemical Company
New York

RECOMMENDATIONS

CABBAGE, BROCCOLI, CAULIFLOWER: Downy Mildew, Leaf Spot.

CARROTS: Cercospora Blight, Macrosporium Blight.

CELERY: Early Blight, Late Blight, and Bacterial Blight. Dust should be applied at weekly intervals in the seedbed and every 7 to 10 days in the field.

PEARS: Fire blight — Apply 13-25 lbs. per acre. Start dusting at 10% bloom and repeat at 5-day intervals until late bloom is over. Should rain occur during this period, shorten interval to 3 days. A dusting should be made within 72 hours prior to expected rain. Do not apply to Anjou or Comice varieties.

PEPPERS: Anthracnose, Cercospora leaf spot.

STRAWBERRIES: Leaf spot (Mycosphaerella).

TOMATOES: Late Blight or Septoria Leaf Spot and Early Blight. It is important to dust plants 5 days before, or 5 days after transplanting.

WALNUTS: To control walnut blight apply at the rate of 30-50 lb. per acre. Apply when catkins are fully expanded and again when nutlets are in 10% bloom. Repeat at 5-day intervals if infective conditions persist.

50 LB. NET

BURN EMPTY BAG — KEEP OUT OF SMOKE