

[476-121]



# ROTENONE S-WP

WETTABLE POWDER • FOR PLANT INSECT CONTROL

### Active Ingredients:

Rotenone .....	5%
Other ethyl ether extractives of cube .....	5%
Inert Ingredients .....	90%
	<hr/>
	100%

## CAUTION

### Keep Out of Reach of Children

Harmful if swallowed or absorbed through skin. Avoid prolonged or repeated contact with skin. Wear clean clothing.

### DIRECTIONS FOR MIXING

First prepare a slurry of Rotenone S-WP and water in a separate container, then add to the spray tank. Keep agitator running during filling and spraying operations.

### OBSERVE FOLLOWING PRECAUTIONS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of spray from areas treated. Do not use rotenone when temperature is below 62° F. Do not apply within 24 hours of harvest.

**BURN EMPTY BAG — KEEP OUT OF SMOKE**

### RECOMMENDATIONS

Unless otherwise stated, applications should be made at the first sign of infestation. Repeat applications should be made as field conditions warrant.

Use 3-5 pounds of Rotenone S-WP per acre application in sufficient water to provide uniform and thorough coverage.

**ASPARAGUS** — Asparagus beetle, 12 spotted asparagus beetle. Apply during cutting period.

**BEANS** — Bean leaf beetle, cucumber beetle, Mexican bean beetle.

**BERRIES** (Blackberry, Boysenberry, Dewberry, Loganberry, Raspberry) — Raspberry cane borer, raspberry fruitworm, raspberry sawfly, red-necked cane borer. Apply on foliage just before plants bloom. Repeat in about 2 weeks. For fruitworm, apply 7 days after first blossoms; repeat in 10 and 20 days.

**CURRENT, GOOSEBERRY** — Gooseberry fruitworm, imported currantworm. Apply when webbing is observed and 7-10 days later.

**CRUCIFERS** (Broccoli, Brussels sprouts, cabbage, cauliflower) — Flea beetle, vegetable weevil, cabbage looper, imported cabbage worm, diamond-back moth.

**LETTUCE** — Aphid, cabbage loopers, flea beetles.

**STRAWBERRY** — Flea beetle, spittle bug, strawberry leaf beetle.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.


4 LBS NET

LOT NO. \_\_\_\_\_

USDA Reg. No. 476-121

S-SF-700326-NP

**STAUFFER CHEMICAL COMPANY**  
NEW YORK, NY 10017





# ROTENONE S-WP

WETTABLE POWDER • FOR PLANT INSECT CONTROL

### Active Ingredients:

Rotenone	5%
Other ethyl ether extractives of cube	5%
Inert Ingredients	90%
	-----
	100%

### RECOMMENDATIONS

Unless otherwise stated, applications should be made at the first sign of infestation. Repeat applications should be made as field conditions warrant.

For best results, use 1/2 to 1 lb. of Rotenone S-WP per acre in sufficient water to provide uniform and thorough coverage.

**ASPARAGUS** - Asparagus beetle, 1-spotted asparagus beetle. Apply during cutting period.

**BEANS** - Bean leaf beetle, cucumber beetle, Mexican bean beetle.

**BERRIES** (Blackberry, Boysenberry, Dewberry, Loganberry, Raspberry) - Raspberry cane borer, raspberry fruitworm, raspberry sawfly, red necked cane borer. Apply on foliage just before plants bloom. Repeat in about 2 weeks. For fruitworm, apply 7 days after first blossoms; repeat in 10 and 20 days.

**CURRENT, GOOSEBERRY** - Gooseberry fruit worm, imported currantworm. Apply when webbing is observed and 7-10 days later.

**CRUCIFERS** (Broccoli, Brussels sprouts, cabbage, cauliflower) - Flea beetle, vegetable weevil, cabbage looper, imported cabbage worm, diamond-back moth.

**LETTUCE** - Aphid, cabbage loopers, flea beetles.

**STRAWBERRY** - Flea beetle, spittle bug, strawberry leaf beetle.

## CAUTION

### Keep Out of Reach of Children

Do not swallow or inhale dust. Avoid contact with skin and eyes. Wash hands after use. Avoid contact with clothing.

### DIRECTIONS FOR MIXING

First prepare a slurry of Rotenone S-WP and water in a separate container, then add to the spray tank. Keep agitator running during filling and spraying operations.

### OBSERVE FOLLOWING PRECAUTIONS

This product is toxic to fish. Keep out of streams and ponds. Do not apply when weather conditions favor drift spray. Do not apply to areas treated with other insecticides when temperature is below 60°F. Do not apply within 24 hours of harvest.

BURN EMPTY BAG - KEEP OUT OF SMOKE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and, or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

4 LBS NET

LOT NO. \_\_\_\_\_

USDA Reg. No. 476-121

S-SF-700326-NP

**STAUFFER CHEMICAL COMPANY**  
NEW YORK, NY 10017

ACCEPTED  
6-3-70  
476-121