

ACCEPTED

FEB 12 1964

UNDER THE FEDERAL FOOD,
DRUG, AND COSMETIC ACT,
FUNGICIDE AND RODENTICIDE AND
FOR ECONOMIC POISON AND HERB-
ICIDE UNDER NO. 272, 747 SUBJECT
TO ATTACHED COMMENTS.

... ..
... ..
... ..
... ..

USE

... ..
... ..
... ..
... ..

USE

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

USE


... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..

... ..
... ..
... ..
... ..



NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

CITRUS: Citrus red and citrus rust mites — Apply 80 lbs. per acre; (1/2 to 1-1/2 lbs. per tree, depending on size for citrus rust mite in Florida). Citrus flat and Yuma spider mites — Apply 100 lbs. per acre. Citrus thrips — Apply 100 lbs. per acre in three applications; mid-March, 1 fall and 2 4 weeks later. Do not apply within 3 to 6 weeks of an oil spray.

COFFEES (MINIPEA VARIETIES): Rust, powdery mildew — Apply 15 to 20 lbs. per acre in 4 applications when the disease is first observed; 12-15 days afterwards; 10-14 days thereafter. Caution: Sulfur is extremely phytotoxic to foliage of Coffea and most other species and should not be used on these varieties.

TEA: Rust, powdery mildew, citrus red, citrus rust mites, citrus thrips, spider mites — Apply 100 lbs. per acre. Do not apply within 3 to 6 weeks of an oil spray.

PEACH: Rust, powdery mildew, citrus red, citrus rust mites, citrus thrips, spider mites — Apply 100 lbs. per acre. Do not apply within 3 to 6 weeks of an oil spray.

PLUM: Rust, powdery mildew, citrus red, citrus rust mites, citrus thrips, spider mites — Apply 100 lbs. per acre. Do not apply within 3 to 6 weeks of an oil spray.

WILDCROPS

ALMOND: Rust, powdery mildew — Apply 10-25 lbs. per acre. Caution: Sulfur may cause petal burn if applied too late.

BEANS: Rust, powdery mildew, spider mites — Apply 15-30 lbs. per acre. For rust control, start applications when the first true leaves form and continue at weekly intervals until late in the season. May cause injury to some varieties of beans.

COTTON: Rust, powdery mildew — Apply 25-50 lbs. per acre. For rust control, start applications when the first true leaves form and continue at weekly intervals until bloom time.

COTTON: (Soybean and soybean rust only) — Powdery mildew, spider mites — Apply 100 lbs. per acre.

PEACH: Rust, powdery mildew — Apply 15-20 lbs. per acre.

PEACH: Rust, powdery mildew — Apply 15-20 lbs. per acre.

PEACH: Rust, powdery mildew — Apply 15-20 lbs. per acre.


GENERAL: Make full coverage applications on both sides of leaves. Keep new growth covered. Chrysanthemum: For early control, do not apply when flowering. Hollyhock: Rust: Do not apply when in flower. Caution: Do not apply when the weather is hot and dry.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

U.S. PATENT OFFICE

REGISTERED TRADEMARK

U.S. PATENT OFFICE



Cameraman's Note
Poor Copy




SWAN[®]

BRAND

DUSTING SULFUR

AN AGRICULTURAL INSECTICIDE and FUNGICIDE



Active Ingredient:

Sulfur

Inert Ingredients

97.5%

2.5%

100%

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground power equipment or aircraft. Except as otherwise shown, apply at the first sign of infection and repeat at 7-10 day intervals as needed to maintain control.

RECOMMENDATIONS

FRUIT CROPS

Apples: Scab, powdery mildew -- Apply 30-35 lbs. per acre in the green tip to pink bud stage. Use on sulfur tolerant varieties only.

Bushberries: Powdery mildew -- Apply 15-20 lbs. per acre.

Cherry: Blossom blight, brown rot, powdery mildew -- Apply 50 lbs. per acre in the popcorn, full bloom and petal fall stages. Leaf spot, powdery mildew -- Apply 50 lbs. per acre at petal fall, shuck fall and 2 weeks after shuck-fall.

Citrus: Six spotted and citrus rust mites -- Apply 80 lbs. per acre, citrus flat and Yuma spider mites -- Apply 100 lbs. per acre, citrus thrips -- Apply 100 lbs. per acre in three applications, mid March, petal fall and 2-4 weeks later. Avoid applications during or preceding high temperatures. Do not apply within 2 months of an oil spray.

Grapes: Rust, powdery mildew -- Apply 5-10 lbs. per acre in 4 applications when shoots are 6" long, shoots 12-15" long; 14 days afterwards; 14 days thereafter to harvest.

Plum-Prune: Brown rot, leaf spot, prune rust -- Apply 35-50 lbs. per acre starting 5 weeks before harvest and continuing at weekly intervals.

Strawberries: Powdery mildew -- Apply 10-20 lbs. per acre.

VEGETABLE & FIELD CROPS

Alfalfa: Spider mites -- Apply 15-25 lbs. per acre. Caution: sulfur may cause petal burn to open flowers.

Beans: Powdery mildew, rust, spider mites -- Apply 15-30 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. Caution: May cause injury to some varieties of beans.

Carrots: Petrobia mite -- Apply 30-35 lbs. per acre.

Clover: Powdery mildew, spider mites -- Apply 15-25 lbs. per acre.

Cotton: Atlantic mite, cotton rust -- Apply 15-30 lbs. per acre.

Grass Seed Crops: Timothy mite -- Apply 30 lbs. per acre.

Melon: Honeydew and sulfur resistant melons only -- Powdery mildew, spider mites -- apply 20-25 lbs. per acre.

Onions: Petrobia mite -- Apply 20-30 lbs. per acre during warm weather.

Peas: Powdery mildew -- Apply 15-20 lbs. per acre.

Sugar Beets: Spider mite -- Apply 30-35 lbs. per acre.

Tomatoes: Tomato russet mite -- Apply 15-40 lbs. per acre.

Turnips and Rutabagas: Powdery mildew -- Apply 15-20 lbs. per acre.

ORNAMENTALS:

Make full coverage applications on all sides of branches. Keep new growth protected.

Chrysanthemum: Powdery mildew, rust -- Apply 15-20 lbs. per acre.

Hollyhock: Rust.

Rose, Perennials, Shrubs: Black spot, powdery mildew, rust. Caution: some burning may occur when used during warm weather.

CAUTION Keep Out of Reach of Children

USE PRECAUTIONS

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on apricots, Anjou and comice pears, cranberries, cucurbits (cucumbers, squash and some varieties of melons), filberts, spinach, tung trees, walnuts and other sensitive plants.

Except on cotton, do not apply when shade temperatures exceed or are likely to exceed 90° F.

Sulfur may burn foliage and fruit during periods of high temperatures and under certain climatic conditions.

Do not use sulfur within 30 days of an oil spray (60 days on citrus).

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

BURN EMPTY BAG—KEEP OUT OF SMOKE

50 LB. NET

U.S.D.A. Reg. No. 476-172

122-122-100

STAUFFER CHEMICAL COMPANY NEW YORK, NY 10017