



MOUSE KILLER

ACTIVE INGREDIENTS:

Warfarin 1² (α-acetonylbenzyl)-4-hydroxycoumarin) 0.025%

N-(2-Quinolyl)-5-sulfanilamide (Sulfaquinoxaline) 0.025%

INERT INGREDIENTS: 99.950%

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Read cautions and directions on back before use

EPA Registration No. 475-183-AA

NET WT. 2 OZ. (56 Gm.)

HOW DOES BLACK FLAG KILL? Contains Prolin which is Warfarin + Sulfaquinoxaline. When eaten by mice over a period of several days, the killing ingredient Warfarin causes internal bleeding and eventual death. It is a progressive action. That's why it is an absolute necessity to feed continuously until all signs of feeding are gone.

Sulfaquinoxaline is an antibacterial agent that inhibits the growth of bacteria that produces Vitamin K in rodents.

NEW SEALED PLASTIC POUCHES REMAIN UNOPENED so placement is neater. Black Flag's deadly ingredients are locked inside so hands never touch bait.

NO MESS, NO SPILL. Unlike other rodenticides there's no mess or spilling during placement. Packaging also keeps bait fresh and attractive until rodents nibble through.

EASIER-TO-PLACE. Because plastic pouches are flexible, you can place them in hard-to-reach areas like behind appliances, in corners, around pipes.

EPA Est. 9562-TX-1

← FRONT OF POUCH

DIRECTIONS

FOR MICE, USE AN UNOPENED POUCH near burrows and runways. Protect from children, pets or livestock with boxes or cages where necessary. Use at 8 to 12 foot intervals.

FOR BEST RESULTS: Check bait daily.

FOR MICE, bait continuously for not less than 15 days. Continue bait until all signs of feeding have stopped.

IMPORTANT: If bait becomes damp, mouldy or stale, replace with fresh pouch immediately.

CAUTION: Protect children, pets, wildlife and domestic animals from the bait. If swallowed by humans, domestic animals or pets, call a physician or veterinarian immediately. The product may reduce the clotting ability of blood and cause internal bleeding. Antidote—Vitamin K.

Baits should be placed in areas not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes. Do not allow treated bait to be exposed on bare ground. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ADDITIONAL DIRECTIONS: Pouches not touched after a week of checking should be moved to more active locations. It is important to feed bait continuously for as long as there are signs of feeding. Interrupted feeding may allow affected rodents to recover.

← BACK OF POUCH

Distributed by **BOYLE-MIDWAY INC., NEW YORK, N.Y. 10017**

Boyle-Midway Div.

Made in U.S.A.

BFMK PI

