

Stauffer
CHEMICALS

ANCHOR
VELVET FLOWERS
OF SULFUR

BRAND OF
SUBLIMED SULFUR N. F.

ACTIVE INGREDIENT: SULFUR.. 99.8%

INERT INGREDIENTS: 0.2%

ANCHOR is a Registered Trademark of Stauffer Chemical Co. 100.0%

STAUFFER CHEMICAL CO.

NEW YORK

50
LB.NET

ANCHOR VELVET FLOWERS OF SULFUR BRAND OF SUBLIMED SULFUR N.F.

CAUTION

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply to epri-cots, comice pears, cranberries, cucurbits (or zucchini, cantaloupes, melons, squash), wal-nuts, or to other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90° F.

Do not use sulfur with oil sprays or within 30 days of an oil spray.

Do not use sulfur on any crop unless local use has proved safe.

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5 to 10 day intervals, depending on severity of infection, but observe urea limits given for specific crops. Apply at the rate of 30 to 40 lb. unless otherwise specified.

RECOMMENDATIONS

FRUIT CROPS

APPLES: Scab, powdery mildew—Apply 30-40 lbs. per acre in the green tip to pink bud stage. Use on sulfur tolerant varieties only.

BUSHBERRIES: Powdery mildew—Apply 15-20 lbs. per acre.

CHERRY: Blossom blight, brown rot, powdery mildew—Apply 50 lbs. per acre in the popcorn, full bloom and petal fall stages; leaf spot, powdery mildew—Apply 50 lbs. per acre at petal fall, shoot fall and 2 weeks after shoot fall.

CITRUS: Six-spotted and citrus rust mites—Apply 80 lbs. per acre (1/2 to 1 1/2 lbs. per tree, depending on size, for citrus rust mite in Florida); Citrus Pith and Yema spider mites—Apply 100 lbs. per acre; citrus thrips—Apply 40 lbs. per acre in three applications: mid-March, petal fall and 2-4 weeks later. Do not apply within 3 to 6 weeks of an oil spray.

GRAPES (VITIFERA VARIETIES): Rust, powdery mildew—Apply 15 to 20 lbs. per acre in 4 applications: when shoots are 6" long; shoots 12-15" long; 14 days after-wards; 10-14 days thereafter to harvest. Caution: Sulfur is extremely phytotoxic to foliage of Concord and most other Isabella type grape varieties. Do not be used on these varieties.

PEACH, NECTARINE: Brown rot, Coryneum blight, rust, peach silver mite, powdery mildew—Apply 50 to 60 lbs. per acre from popcorn stage to harvest.

PLUM, PRUNE: Brown rot, leaf spot, prunella rust—Apply 30-40 lbs. per acre starting 5 weeks before harvest and continuing at weekly intervals.

STRAWBERRIES: Powdery mildew—Apply 15-30 lbs. per acre.

VEGETABLE & FIELD CROPS

ALFALFA: Spider mites—Apply 15-25 lbs. per acre. Caution: Sulfur may cause petal burn to open flowers.

BEANS: Powdery mildew, rust, spider mites—Apply 15-20 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue weekly intervals until blossom time. Caution: May cause injury to some varieties of beans.

COTTON: Atlantic and cotton rust—Apply 25-30 lbs. per acre.

MILKWEED: (Blue, yellow and white varieties only)—Powdery mildew, spider mites—Apply 25-35 lbs. per acre.

PEAS: Powdery mildew—Apply 15-20 lbs. per acre.

PEANUTS: Rust, leaf spot—Apply 15-30 lbs. per acre.

TOMATOES: Tomato leaf blight—Apply 15-20 lbs. per acre.

ANCHOR VELVET FLOWERS OF SULFUR

ANCHOR VELVET FLOWERS OF SULFUR is a registered trademark of Stauffer Chemical Company. Keep away from children. Do not use in confined spaces. Do not use in areas where there is a fire hazard. Do not use in areas where there is a risk of explosion. Do not use in areas where there is a risk of fire. Do not use in areas where there is a risk of injury to persons or animals. Do not use in areas where there is a risk of damage to property. Do not use in areas where there is a risk of damage to crops. Do not use in areas where there is a risk of damage to the environment. Do not use in areas where there is a risk of damage to the ozone layer. Do not use in areas where there is a risk of damage to the climate. Do not use in areas where there is a risk of damage to the soil. Do not use in areas where there is a risk of damage to the water supply. Do not use in areas where there is a risk of damage to the atmosphere. Do not use in areas where there is a risk of damage to the biosphere. Do not use in areas where there is a risk of damage to the hydrosphere. Do not use in areas where there is a risk of damage to the lithosphere. Do not use in areas where there is a risk of damage to the geosphere. Do not use in areas where there is a risk of damage to the cryosphere. Do not use in areas where there is a risk of damage to the atmosphere. Do not use in areas where there is a risk of damage to the hydrosphere. Do not use in areas where there is a risk of damage to the lithosphere. Do not use in areas where there is a risk of damage to the geosphere. Do not use in areas where there is a risk of damage to the cryosphere.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

APPLES: Scab, powdery mildew — Apply 30-40 lbs. per acre in the green tip to pink bud stage. Use on sulfur tolerant varieties only.

BUSHBERRIES: Powdery mildew — Apply 15-20 lbs. per acre.

CHERRY: Blossom blight, brown rot, powdery mildew — Apply 50 lbs. per acre in the popcorn, full bloom and petal fall stages; leaf spot, powdery mildew — Apply 50 lbs. per acre at petal fall, shuck-fall and 2 weeks after shuck-fall

CITRUS: Six-spotted and citrus rust mites — Apply 80 lbs. per acre; (1/2 to 1-1/2 lbs. per tree, depending on size, for citrus rust mite in Florida). Citrus flat and Yuma spider mites — Apply 100 lbs. per acre; citrus thrips — Apply 100 lbs. per acre in three applications; mid-March, petal fall and 2-4 weeks later. Do not apply within 3 to 6 weeks of an oil spray.

GRAPES (VINIFERA VARIETIES): Rust, powdery mildew — Apply 15 to 20 lbs. per acre in 4 applications: when shoots are 6" long; shoots 12-15" long; 14 days afterwards; 10-14 days thereafter to harvest. **Caution:** Sulfur is extremely phytotoxic to foliage of Concord and most other *Vitis* type grapes and should not be used on these varieties.

PEACH, NECTARINE: Brown rot, Coryneum blight, rust, peach silver mite, powdery mildew — Apply 50 to 60 lbs. per acre from popcorn stage to harvest.

PLUM, PRUNE: Brown rot, leaf spot, prune rust — Apply 35-50 lbs. per acre starting 5 weeks before harvest and continuing at weekly intervals.

STRAWBERRIES: Powdery mildew — Apply 15-30 lbs. per acre.

VEGETABLE & FIELD CROPS

ALFALFA: Spider mites — Apply 15-25 lbs. per acre. **Caution:** Sulfur may cause petal burn to open flowers.

BEANS: Powdery mildew, rust, spider mites — Apply 15-30 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. **Caution:** May cause injury to some varieties of beans.

COTTON: Atlantic mite, cotton rust — Apply 25-30 lbs. per acre.

MELON: (Honeydew and sulfur-resistant melons only) — Powdery mildew, spider mites — Apply 20-25 lbs. per acre.

PEAS: Powdery mildew — Apply 15-20 lbs. per acre.

PEANUTS: Rust, leaf spot — Apply 15-30 lbs. per acre.

TOMATOES: Tomato russet mite — Apply 15-30 lbs. per acre.

CHRYSANTHEMUMS: Make full coverage applications on both sides of leaves. Keep new growth protected. **Chrysanthemum:** Powdery mildew, rust. Do not apply when flowering. **Hydrangea:** Rust. **Rose:** Black spot, powdery mildew, rust. **Caution:** Some burning may occur when used during warm weather.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

CROWN 90

WETTABLE

SULFUR

FOR SPRAYING FRUITS,
VEGETABLES, ORNAMENTALS

ACTIVE INGREDIENT: SULFUR 90%

INERT INGREDIENTS 10%

100%

• CROWN is a registered trademark of Stauffer Chemical Company.

FINENESS: not less than 93% passing 325 mesh

CAUTION

EPA Reg. No. 476-1125-AA / KEEP OUT OF REACH OF CHILDREN

Made in U.S.A. by

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880



CROWN 90
SULFUR

1315