



STANDARD LEAD ARSENATE

AN AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENT:	
Standard Lead Arsenate(PbHAsO ₄)	95%
INERT INGREDIENTS	5%
	100%
Lead expressed as Metallic	56.7%
Arsenic expressed as Metallic	20.5%
Arsenic in water soluble form expressed as Metallic, not more than	0.5%

To protect bees, do not apply when crops are in bloom. Remove or cover hives if application is necessary during bloom.

Do not use this product on feed or forage to be fed dairy animals or to livestock being finished for slaughter.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number and rate of applications needed will vary with local conditions.



ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonfuls of Epsom Salts or Milk of Magnesia in water, and plenty of milk and water. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY!

DANGER!

Poisonous if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid breathing dust or spray mist. Do not contaminate feed and food.

Keep Out of Reach of Children and Domestic Animals.

OBSERVE THE FOLLOWING USE PRECAUTIONS

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Toxic to fish and wildlife. Do not contaminate any body of water or apply to any area not specified on the label.

NOTICE: Stauffer Chemical Company makes no warranty of merchantability or any other express or implied warranty concerning this material. It shall not be held responsible for personal injury, property damage, or other loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability resulting from such handling, storage or use.

DIRECTIONS FOR USE

Measure the specified amount of this product then wash through filler screen in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-pump spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants. Emulsifiable concentrates should always be thoroughly emulsified with at least half of total water before adding this product.

RECOMMENDATIONS

Rates are given in pounds of Standard Lead Arsenate per 100 gallons of water in full coverage sprays.

CHERRIES: Cherry Fruit Fly, Black Cherry Fly — Use 2½ to 3 lbs. plus 2½ to 3 lbs. hydrated lime. Begin all treatments when first adults are trapped. Make 2 or 3 applications at 10-14 day intervals.

Pear Slug — Use 2 lbs. plus 2 lbs. hydrated lime making 1 or 2 applications; the first about 20 days after bloom. Do not apply on cherries within 30 days of harvest. (May be applied up to 14 days of harvest on processing cherries if dosage rate limited to 24 lbs. per acre.) Dosage rate for English Morello variety should be reduced to 1 lbs. per 100 gal. water.

PEARS: Codling Moth — Use 3 lbs. Make 2 to 4 cover sprays beginning with calyx.

Pear Leaf Worm — Use 2 lbs. at petal fall

Pear Slug — Use 2 lbs. about two weeks after bloom.

Pear Curculio — Use 3 lbs. at petal fall and 10 days later. Do not apply within 30 days of harvest. Remove excess residue at time of harvest. DO NOT GRAZE LIVESTOCK IN TREATED GROVES.

USDA Reg. No. 476-374 S-SF-660511-WC

BURN BAG WHEN EMPTY — BURY ASHES

4 LBS. NET

STAUFFER CHEMICAL CO.

NEW YORK