

RECOMMENDATIONS

SPECIAL CROPS & USES


ANTS - THE USE OF CHLORDANE 8-EC FOR ANT CONTROL IS LIMITED TO SOIL TREATMENT AT PLANTING, PREPLANTING OR TRANSPLANTING TIME FOR CROPS LISTED ON THIS LABEL; GOLF COURSES AND GREENHOUSES. USE AT THE RATE OF 1 TO 1-1/2 PINTS IN 100 GALLONS OF WATER PER ACRE. APPLY DIRECTLY ON ANTS, ALONG THEIR TRAILS, AND DOUSE HILLS OR NESTS.

GRASSHOPPER CONTROL - (ADJACENT TO BEANS, COLE CROPS LISTED ON THIS LABEL, CORN, PEAS, TOMATOES, STRAWBERRIES AND SMALL GRAINS) - FOR LIMITED USE IN HATCHING AREAS, IN FIELD MARGINS AND FENCE ROWS TO CONTROL YOUNG HOPPERS, USE 1/2 PINT PER 100 GALLON OF WATER PER ACRE; FOR MATURE FORMS, USE 1 PINT PER 100 GALLON PER ACRE. WHEN HOPPERS ARE MIGRATING FROM FIELD TO FIELD, MAKE BARRIER APPLICATION TO THE FIELD MARGINS, OF CROPS LISTED, BORDERING ADVANCING HOPPERS.

LAWNS, ORNAMENTAL TURF - ANTS, CHINCH BUGS, JAPANESE BEETLE LARVAE, WEBWORMS, WHITE GRUBS (JUNE BEETLE LARVAE), FOR ANTS, CHINCH BUGS AND WEBWORMS, USE 2 TO 2-1/2 QUARTS PER ACRE. (1-1/2 OUNCES PER 1,000 SQUARE FEET) IN SUFFICIENT WATER FOR THOROUGH COVERAGE AT FIRST SIGN OF INFESTATION. REPEAT IN 14 DAYS IF NEEDED. FOR WHITE GRUBS AND JAPANESE BEETLE LARVAE, USE 5 QUARTS PER ACRE (3 OUNCES PER 1,000 SQUARE FEET) IN SUFFICIENT WATER FOR COVERAGE TO INFESTED AREAS AND AREAS TO BE PROTECTED. FOR CRABGRASS - APPLY 1-1/2 PINTS IN SUFFICIENT WATER TO THOROUGHLY COVER 1,000 SQUARE FEET. TIMING OF APPLICATION IS IMPORTANT AND MUST BE MADE BEFORE CRABGRASS SEED GERMINATES. CONSULT EXPERIMENT STATION OR EXTENSION SERVICE SPECIALIST AS APPLICATION IS MOST EFFECTIVE IN AREAS IN WHICH DEFINITE SEASONAL GERMINATION OCCURS. IN GENERAL, APPLICATIONS MAY BE MADE FROM VERY EARLY SPRING UNTIL 7 TO 14 DAYS PRIOR TO NORMAL GERMINATION TIME. DO NOT APPLY TO NEWLY SEEDED LAWNS.

CAUTION: DO NOT ALLOW CHILDREN OR PETS ON TREATED LAWNS OR TURF UNTIL INSECTICIDE IS WASHED INTO THE SOIL BY WATERING AND GRASS IS COMPLETELY DRY.

TERMITES - ADD 2 TO 4 QUARTS IN 100 GALLONS WATER. APPLY 1/2 GALLON OF THE SOLUTION PER LINEAR FOOT IN TRENCHES NOT EXCEEDING 15 INCHES IN DEPTH AND 8 INCHES WIDE. FOR BUILDINGS WITH DEEP FOOTINGS, DIG DEEPER TRENCHES. FOR TRENCHES EXCEEDING 15 TO 30 INCHES, APPLY 1 GALLON OF SOLUTION PER LINEAR FOOT. DO NOT DIG TRENCHES BELOW THE LEVEL OF THE FOOTING. INSPECT AREA YEARLY TO DETERMINE POSSIBLE REINFESTATION.



PRE-PLANT SOIL TREATMENT

BROADCAST SOIL APPLICATION

MIX THE REQUIRED AMOUNTS OF CHLORDANE 8-EC IN SUFFICIENT WATER FOR UNIFORM COVERAGE OF THE SOIL SURFACE. APPLICATION SHOULD BE FOLLOWED PROMPTLY BY TILLAGE, OR AS OTHERWISE SPECIFIED FOR SPECIFIC USES.

CORN - CORN ROOTWORM, WHITE-FRINGED BEETLE LARVAE. USE 1 QUART PER ACRE IN 100 GALLONS OF WATER.

CUTWORMS - BEANS, CORN, EGGPLANT, LETTUCE, ONIONS, PEPPERS, POTATOES AND TOMATOES. USE 1 PINT PER 100 GALLONS WATER (150 GALLONS PER ACRE) BEFORE PLANTING. APPLY EVENLY OVER SOIL SURFACE AND DO NOT DISTURB SOIL FOR 3 TO 5 DAYS.

MOLE-CRICKETS - BROCCOLI, CABBAGE, CAULIFLOWER, LETTUCE, STRAWBERRIES AND TOMATOES. APPLY 1 TO 2 PINTS PER ACRE EVENLY OVER SOIL SURFACE BEFORE SEEDING OR TRANSPLANTING, USING SUFFICIENT WATER TO INSURE COMPLETE COVERAGE.

WHITE GRUBS - POTATOES, STRAWBERRIES AND SUGAR BEETS. APPLY 1-1/2 TO 2-1/2 QUARTS IN 100 GALLONS WATER PER ACRE AS A PRE-PLANT TREATMENT TO SOIL SURFACE. DISK INTO TOP 3 OR 4 INCHES OF SOIL.

WIREWORMS - BEANS, BEETS, BROCCOLI, CABBAGE, CORN, CAULIFLOWER, ONIONS, POTATOES, STRAWBERRIES, SUGAR BEETS, SWEET POTATOES AND TURNIPS. MAKE PRE-PLANTING TREATMENT TO SOIL SURFACE APPLYING 1-1/2 TO 2-1/2 QUARTS IN 100 GALLONS WATER PER ACRE. DISK INTO TOP 4 TO 6 INCHES OF SOIL. USE HIGHER DOSAGE ON MUCK SOILS.

WIREWORMS-POTATOES (PACIFIC NORTHWEST) - MAKE PREPLANT TREATMENT TO THE SOIL SURFACE APPLYING 1 GALLON IN 100 GALLONS WATER PER ACRE. DISC INTO TOP 6 TO 9 INCHES OF SOIL.

NOTICE: STAUFFER CHEMICAL COMPANY MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND/OR FITNESS FOR ANY PARTICULAR PURPOSE, CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON STAUFFER'S LABEL.

