


476-655



CAPTAN 10 DUST

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS	
CAPTAN: N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide	10%
INERT INGREDIENTS:	90%
	100%

Not licensed for use or export outside of the United States, its territories and possessions, Canada or Mexico.

CAUTION Keep Out of Reach of Children

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST.
 AVOID CONTACT WITH SKIN.
 AVOID CONTAMINATION OF FEED AND FOOD.
 STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT.

USE PRECAUTIONS

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of dust from areas treated.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5-10 day intervals as needed to maintain control.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Made in U.S.A. — Stauffer Chemical Co., New York

U.S.D.A. Reg. No. 476-655

S-SF-690213-WC

50 LB. NET

RECOMMENDATIONS

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight — Use 40 lbs. per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLE: Scab, Frogeye Leaf Spot — 30 to 40 lbs. per acre as required to maintain adequate coverage in the pre-bloom and blossom applications.

APRICOTS: Brown Rot, Jacket Rot — 30 to 40 lbs. per acre. Apply at red bud stage and repeat at 75 per cent petal fall.

CHERRY: Brown Rot, Leaf Spot — 30 to 40 lbs. per acre. Apply at popcorn, blossom, and petal fall stages, continue at 10 to 14-day intervals, including pre-harvest, if necessary.

GRAPES: Mold Control on Fruit on Drying Trays in the Field (*Penicillium* spp., *Aspergillus* spp., *Botrytis cinerea*, *Rhizopus*, spp.) Apply 10 lbs. per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply this dust by hand or ground dusters only.

GRAPES: Botrytis Rot (Bunch Rot) — 40 to 50 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically up to 3 cover applications before bunches close.

PEACH, NECTARINES: Brown Rot, Peach Scab — 40 to 50 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.

RASPBERRIES: Fruit Rot — 25 lbs. per acre. Make applications 3 to 5 days before harvest starts, at mid-harvest and 8 to 10 days after mid harvest.

STRAWBERRIES: Botrytis Rot — 30 to 38 lbs. per acre beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking, if conditions favor fruit rot.

BEANS, PEAS: Soil Treatment — Damp-off (Beans only), **Root Rot** — Work into top 3 or 4 inches of soil before planting. Apply 40 to 50 lbs. per acre.

CELERY: (Plant bed only) — Damp-off, Pink Rot — Apply 35 to 40 lbs. per acre when all seeds have germinated. Repeat if necessary. Repeat at 5 to 10-day intervals until danger of infection is past.

CUCURBITS (Cantaloupe, Cucumber, Melon, Squash): Angular Leaf Spot, Anthracnose, Downy Mildew — Make application when rotifers begin to form or at the first sign of disease. Apply at the rate of 20 to 30 lbs. per acre on young plants increasing to 30 to 38 lbs. per acre on mature plants.

PLUMS, PRUNES: Brown Rot — 40 to 50 lbs. per acre. Apply at popcorn, full bloom, petal fall and 2 covers as necessary. Thorough coverage is essential for maximum control.

POTATOES: Early and Late Blight — Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens. Apply 30 to 35 lbs. per acre on young plants. Use 40 lbs. per acre on mature plants.

POTATO SEED PIECE TREATMENT: Damp-off, Seed Rot, Brown Eye Disease, Verticillium Wilt — Use at rate of 10 ozs. per bushel of seed pieces (16 oz. or 1 lb. to 100 lbs. of cut seed piece). Apply in convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates two days before bagging.

TOMATOES: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot — Apply 20 to 30 lbs. per acre at weekly intervals on young plants; 40 to 50 lbs. per acre on mature plants.

SOIL AND GREENHOUSE BENCH TREATMENT: 10 lbs. per 1,000 sq. ft. (60 lbs. per acre). Work into top 3 or 4 inches of soil before planting. For damp off and root rot diseases of seedlings and transplants. Effective on flowers, lawn seedbeds, and vegetables such as beans, peas, cabbage, eggplant, pepper and tomato.

ORNAMENTALS: Rose, Black Spot; Carnation, Rust; Leaf Spot; Chrysanthemum, Flower Blight — Apply thoroughly prior to first sign of disease. Repeat applications at 7 to 10 day intervals. Shorten interval during periods of frequent rain and heavy dews.

BURN EMPTY BAG—KEEP OUT OF SMOKE