

Stauffer

CAPTAN 10 DUST

AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENTS:

CAPTAN: N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide	9.46%
Related derivatives	0.54%
INERT INGREDIENTS:	90.00%
	100.0%

CAUTION

Keep Out of Reach of Children

HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST.

AVOID CONTACT WITH SKIN.

AVOID CONTAMINATION OF FEED AND FOOD.

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT.

USE PRECAUTIONS

In order that pesticide residues on food and forage crops will not exceed tolerances established by the Environmental Protection Agency and/or Food and Drug Administration, use only at recommended rates and intervals.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of dust from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing of applications will vary with local conditions.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5-10 day intervals as needed to maintain control.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE Stauffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties; THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which damages are claimed.

EPA Reg. No. 476-655-AA

A-1

EPA Est. 476-CA-1

50 LB. NET

This product is sold only for uses stated on the label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: U.S.A.

Made in U.S.A. by

STAUFFER CHEMICAL COMPANY WESTPORT, CT 06880

RECOMMENDATIONS

ALMONDS: Brown Rot, Shot Hole, Scab, Leaf Blight—Use 40 lbs. per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

APPLE: Scab, Frogeye Leaf Spot—30 to 40 lbs. per acre as required to maintain adequate coverage in the pre-bloom and blossom applications.

APRICOTS: Brown Rot, Jacket Rot—30 to 40 lbs. per acre. Apply at red bud stage and repeat at 75 per cent petal fall.

CHERRY: Brown Rot, Leaf Spot—30 to 40 lbs. per acre. Apply at popcorn, blossom, and petal fall stages. Continue at 10 to 14-day intervals, including pre-harvest, if necessary.

GRAPES: Mold Control on Fruit on Drying Trays in the Field (*Penicillium* spp., *Aspergillus* spp., *Botrytis cinerea*, *Rhizopus*, spp.)—Apply 10 lbs. per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply this dust by hand or ground dusters only.

GRAPES: Botrytis Rot (Bunch Rot)—40 to 50 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically up to 3 cover applications before bunches close.

PEACH, NECTARINES: Brown Rot, Peach Scab—40 to 50 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.

RASPBERRIES: Fruit Rot—25 lbs. per acre. Make applications 3 to 5 days before harvest starts, at mid-harvest and 8 to 10 days after mid-harvest.

STRAWBERRIES: Botrytis Rot—30 to 38 lbs. per acre beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking if conditions favor fruit rot.

BEANS, PEAS: Soil Treatment—Damp-off (Beans only), Root Rot—Work into top 3 or 4 inches of soil before planting. Apply 40 to 55 lbs. per acre.

CELERY: (Plant bed only)—Damp-off, Pink Rot—Apply 35 to 40 lbs. per acre when all seeds have germinated. Repeat if necessary. Repeat at 5 to 10-day intervals until danger of infection is past.

CUCURBITS (Cantaloupe, Cucumber, Melon, Squash): Angular Leaf Spot, Anthracnose, Downy Mildew—Start application when runners begin to form or at the first sign of disease. Apply at the rate of 20 to 30 lbs. per acre on young plants increasing to 30 to 38 lbs. per acre on mature plants.

PLUMS, PRUNES: Brown Rot—40 to 50 lbs. per acre. Apply at popcorn, full bloom, petal-fall and in covers as necessary. Thorough coverage is essential for maximum control.

POTATOES: Early and Late Blight—Make first application when plants are 2 to 4 inches high. Repeat at 7 to 10-day intervals as long as disease threatens. Apply 30 to 35 lbs. per acre on young plants. Use 40 lbs. per acre on mature plants.

POTATO SEED PIECE TREATMENT: Damp-off, Seed Rot Fungi, Brown Eye Disease, Verticillium Wilt—Apply at the rate of 7½ ounces per bushel cut seed pieces (12 ounces or ¾ pound to 100 pounds of cut seed pieces). Apply in convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed, store treated seed pieces in open crates for two days before bagging.

TOMATOES: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot—Apply 20 to 30 lbs. per acre at weekly intervals on young plants; 40 to 50 lbs. per acre on mature plants.

SOIL AND GREENHOUSE BENCH TREATMENT:—1½ lbs. per 1,000 sq. ft. (60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting. For damp-off and root rot diseases of seedlings and transplants of roses, other shrubs, flowers, lawn seedbeds, and vegetables such as beans, peas, cabbage, eggplant, pepper and tomato.

ORNAMENTALS: Rose, Black Spot; Carnation, Rust, Leaf Spot; Chrysanthemum, Flower Blight—Apply thoroughly prior to or at first sign of disease. Repeat applications at 7 to 10-day intervals. Shorten interval during periods of frequent rain and heavy dews.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides, or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.