

ACCEPTED



# CAPTAN 75 SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE CONTROL — DYE ADDED

ACTIVE INGREDIENT:

Captan

75%

INERT INGREDIENTS

25%

*N*-[(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide]

100%

Not licensed for use in countries where valid patents still exist

## CAUTION

## Keep Out of Reach of Children

CAUSES SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED OR INHALED.

Avoid contact with skin, eyes, or clothing. Avoid breathing dust — wear respirator. Wash thoroughly after handling. Do not store or use near feed or food. In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Wash skin with soap and water.

PROTECT FROM EXCESSIVE HEAT — STORE IN A COOL DRY PLACE  
CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES.

**CAPTAN 75 SEED PROTECTANT** is a microfine wettable powder, especially formulated for the treatment of seed, prior to storage and planting, to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

**COMPATIBILITY:** Captan 75 can be used in combination with most commonly used seed protectants, both insecticides and fungicides as well as fertilizers. Do not use with strongly alkaline materials.

**PLANT SAFETY:** At the recommended dosage Captan 75 is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated, giving a superior result as compared to results from other seed fungicides.

## GENERAL USE PRECAUTIONS

Read all directions before using and use only as specified on this label.

Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of excess treated seeds by burial.

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED  
DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

The Federal Government and some states have issued regulations for the application of pesticides to agricultural crops. Check the regulations for your state.

## DIRECTIONS FOR USE

**SLURRY TREATMENT:** Prepare a slurry by adding a part of the product to a quantity of water equal to the weight of the product. Mix this part of the slurry with the seed to be treated.

## RECOMMENDATIONS

## AMOUNT OF CAPTAN 75 TO USE ON SEEDS

# CAUTION

# Keep Out of Reach of Children

CAUSES SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED OR INHALED.

Avoid contact with skin, eyes, or clothing. Avoid breathing dust — wear respirator. Wash thoroughly after handling. Do not store or use near feed or food. In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Wash skin with soap and water.

PROTECT FROM EXCESSIVE HEAT — STORE IN A COOL DRY PLACE  
CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES.

**CAPTAN 75 SEED PROTECTANT** is a microfine wettable powder especially formulated for the treatment of seed, prior to storage and planting, to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

**COMPATIBILITY:** Captan 75 can be used in combination with most commonly used seed protectants, both insecticides and fungicides as well as fertilizers. Do not use with strongly alkaline materials.

**PLANT SAFETY:** At the recommended dosage Captan 75 is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated, giving a superior result as compared to results from other seed fungicides.

## GENERAL USE PRECAUTIONS

Read all directions before using and use only as specified on this label.

Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of excess treated seeds by burial.

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

**TREATED SEED  
DO NOT USE FOR FEED, FOOD OR OIL PURPOSES**

The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

## DIRECTIONS FOR USE

**SLURRY TREATMENT:** Pre mix the recommended amount of this material with a little water in a bucket and make a thin paste. Stir this paste into the required amount of water for the slurry equipment to be used.

## RECOMMENDATIONS

For control of many seed rots, seedling rots, root rots, fungi and kernel smut of sorghum.

## AMOUNT OF CAPTAN 75 TO USE ON SEEDS

SEED	SLURRY	SEED	SLURRY
ALFALFA, CLOVER, LESPEDEZA, TREFOLI	5 1/2 oz. 100 lb	FLAX	1 1/2 oz. bu
BARLEY, OATS, WHEAT	2 1/2 oz. bu	GRASSES	1 1/2 oz. 100 lb
BEANS (Dry, Snap, Lima), Cowpeas, Peas	1 oz. bu	MUSKMELON, PUMPKIN, SQUASH	
BEETS (Garden)	8 oz. 100 lb	WATERMELON	1 1/2 oz. 100 lb
CANTALOUPE, CUCUMBER	2 oz. 100 lb	ONIONS (Planting)	1 lb. 1 lb
COLE CROPS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower)	1 oz. 100 lb	PEANUTS	4 oz. 100 lb
CORN FIELD	2 1/2 oz. bu	RICE	2 1/2 oz. 100 lb
SWEET	1 1/2 oz. 100 lb	SORGHUM (all types)	1 1/2 oz. bu
COTTON ACID DELINTED	1 1/2 oz. 100 lb	SOYBEANS	1 1/2 oz. bu
REGINNED	3 oz. 100 lb	SPINACH	4 oz. 100 lb
FIZZY	3 oz. 100 lb	TIGAR BEETS (all types)	1 1/2 oz. 100 lb
MARBLETS (Mustard, Radishes, Rape, Turnips)	1 1/2 oz. 100 lb	W. S. CHARD	1 1/2 oz. 100 lb

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

EPA Reg. No. 149-100-01

10 LBS. NET

**STAUFFER CHEMICAL COMPANY**

WESTPORT, CT 06880