



CAPTAN 75 SEED PROTECTANT

SLURRY APPLICATION FOR SEED DISEASE CONTROL — DYE ADDED

CAPTAN 75 SEED PROTECTANT is a microfine wettable powder especially formulated for the treatment of seed, prior to storage and planting, to protect seed from molds and other fungi causing storage loss and to protect seed from seed-borne and soil-borne fungi which cause seed decay and seedling blights.

COMPATIBILITY: Captan 75 can be used in combination with most commonly used seed protectants, both insecticides and fungicides as well as fertilizers. Do not use with strongly alkaline materials.

PLANT SAFETY: At the recommended dosage Captan 75 is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated, giving a superior result as compared to results from other seed fungicides.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SLURRY TREATMENT: Pre-mix the recommended amount of this material with a little water in a bucket and make a thin paste. Stir this paste into the required amount of water for the slurry equipment to be used.

NOTICE — READ CAREFULLY

CONDITIONS OF SALE Stauffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties; **THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY** and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

ACTIVE INGREDIENTS:

CAPTAN: N-(trichloromethyl)thio-1,4-cyclohexene-1,2-dicarboximide 70.92%

Related derivatives 4.08%

INERT INGREDIENTS: 25.00%

100.00%

CAUTION

Keep Out of Reach of Children
CAUSES SKIN AND EYE IRRITATION.
MAY BE HARMFUL IF SWALLOWED OR INHALED.

Avoid contact with skin, eyes, or clothing.
Avoid breathing dust — wear respirator.
Wash thoroughly after handling.

Do not store or use near feed or food.
In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Wash skin with soap and water.

PROTECT FROM EXCESSIVE HEAT
STORE IN A COOL DRY PLACE

GENERAL USE PRECAUTIONS

Read all directions before using and use only as specified on this label.

Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Dispose of excess treated seeds by burial.

IMPORTANT: Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED

DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.

RECOMMENDATIONS

For control of many seed rots, seedling rots, root rots, fungi and kernel smut of sorghum.

AMOUNT OF CAPTAN 75 TO USE ON SEEDS

SEED	SLURRY
Alfalfa, Clover, Lespedeza, Trefoil	5½ oz./100 lb.
Barley, Oats, Wheat	2/3 oz./bu.
Beans (Dry, Snap, Lima), Cowpeas, Peas	1 oz./bu.
Beets (Garden)	8 oz./100 lb.
Cantaloupe, Cucumber	2 oz./100 lb.
Cole Crops (Broccoli, Brussels Sprouts, Cabbage, Cauliflower)	1 oz./100 lb.
Corn — Field	2/3 oz./bu.
Sweet	2½ oz./100 lb.
Cotton — Acid Delinted	2 oz./100 lb.
Reginned	3 oz./100 lb.
Fuzzy	3 oz./100 lb.
Crucifers (Mustard, Radishes, Rape, Turnips)	1 oz./100 lb.
Flax	1½ oz./bu.
Grasses	5½ oz./100 lb.
Muskmelon, Pumpkin, Squash, Watermelon	1½ oz./100 lb.
Onions (Pelleting)	1 lb./1 lb.
Peanuts	4 oz./100 lb.
Rice	2¼ oz./100 lb.
Sorghum (Hull-less)	1¼ oz./bu.
Soybeans	1 oz./bu.
Spinach	4 oz./100 lb.
Sugar Beets — West	4 oz./100 lb.
East	8 oz./100 lb.
Swiss Chard	8 oz./100 lb.

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

EPA Reg. No. 476-676-AA EPA Est. 476-OR-1
50 LBS. NET A-1

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL: Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

This product is sold only for uses stated on the label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: U.S.A.

Made in U.S.A. by

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880