

CAPTAN 7.5 DUST

AN AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT
INERT INGREDIENTS

RECOMMENDATIONS

CAUTION

HARMFUL IF TAKEN INTERNALLY

- AVOID INHALE AND DRINK
- AVOID CONTACT WITH SKIN
- AVOID CONTAMINATION OF FEEDSTUFFS
- DO NOT EAT OR DRINK FOR 2 HOURS AFTER USE

USE PRECAUTIONS

In order that residues on food and feed are safe for use, use only at recommended rates. This product is toxic to fish. Keep away from water. Do not apply when weather conditions favor drift of dust. Do not contaminate water by cleaning of equipment or disposal of wash water. Consult State Dept. of Agriculture for additional information. More needed will vary with conditions.

DIRECTIONS FOR USE

It is a violation of Federal law to apply this product in a manner inconsistent with the directions on the label. Do not apply to water.

STATE OF CALIFORNIA

WARRANTY AND LIABILITY

INHERENT RISKS

LIMITATION OF LIABILITY

© 1975 Stauffer Chemical Company

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

Apply to the following crops and diseases. Do not apply to water. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan. Do not apply to crops that are sensitive to Captan.

50 LB NET WT 22.7 kg



STAUFFER CHEMICAL COMPANY

USE AND DISPOSAL

