

Ass
App
Barr
Barr
Blac
Blac
Buc
Buc
Ca
Ce
Che
Che
Che
Che
It
It
It
It
best
best
Stu
Stu
volu
volu
Do
Do
STUN
this
12
kay
taker
were
BRAUS
cont
pian
After
After
FRC
AIRPL
apply
GROU
these
NOT
or is
and/
mate
factu
USDA

WARNING

DO NOT include directly or indirectly identifiable information (e.g., names, addresses, phone numbers, e-mail addresses, or other identifying information) that are sensitive to a 4(C) 2.4(C) 1.1 or 2.4(C) 1.2.

DO NOT burn sprays in the 24" x 24" x 10" incinerator. Suitable plants are listed on the back of the spray can. Overloading, dirty, or old sprays and sprays with chloroforms are less acceptable.

DO NOT apply to any other type of material or other
describes susceptible to fire.

Excessive amounts of fertilizer, particularly nitrogen, can cause rapid growth of plant growths **DO NOT** use fertilizer on plants that are susceptible to this disease. Do not use fertilizer on plants that are susceptible to damage or killed.

Application should be carried out only where there is no hazard from drift

Vapors from this product may injure susceptible plants growing in the immediate vicinity.

1. The first step is to identify the problem or goal. This involves understanding the current situation and what needs to be achieved.

Because of the difficulty of thoroughly cleaning sprayers and other equipment used, such equipment should not be used for handling and applying other agricultural chemicals.

... were sent out of the State Agricultural Extension Service to the State Agricultural Extension Service for additional information on the timing and rate of application necessary with the local conditions.

WOODY PLANT CONTROL Not a growing crop - along fence lines, drainage ditch banks, right of ways, around farm yards and buildings. It is water-soluble at the rate of 1 to 3 gallons in 100 gallons of water. Applied and water will control the following susceptible plants:

(Contains 1.33 lb. 2,4-D, .66 lb. 2,4,5-T Acid Per Gallon)

Butyl ester of 2,4-dichlorophenoxyacetic acid
Butyl ester of 2,4,5-trichlorophenoxyacetic acid

2,4-dichlorophenoxyacetic acid equivalent
2,4,5-trichlorophenoxyacetic acid equivalent

19.5%
9.5%
71.0%

100.0%
15.5%
7.8%

ACCEPTED

HARMFUL IF SWALLOWED
May cause skin irritation.

Protect From Temperatures Below 25° F.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

DESTROY EMPTY CONTAINER IMMEDIATELY — PERFORATE AND BURN COMMENTS.

DO NOT BURN

GAL. NET

2,4-D BUTYL ESTERS 2,4,5-T 1.33-.66-E

HIGH VOLATILE BRUSH KILLER

EMULSIFIABLE LIQUID • FOR DILUTION WITH OIL OR WATER

(Contains 1.33 lb. 2,4-D, .66 lb. 2,4,5-T Acid Per Gallon)

ACTIVE INGREDIENTS:

Butyl ester of 2,4-dichlorophenoxyacetic acid 19.5%

Butyl ester of 2,4,5-trichlorophenoxyacetic acid 9.5%

INERT INGREDIENTS

2,4-dichlorophenoxyacetic acid equivalent 100.0%

2,4,5-trichlorophenoxyacetic acid equivalent 15.5%

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED

May cause skin irritation.

Avoid contact with eyes, skin, and clothing.

Protect From Temperatures Below 25° F.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

DESTROY EMPTY CONTAINER IMMEDIATELY — PERFORATE AND BURY COMMENTS.

DO NOT BURN

ACCEPTED

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 779-733 SUBJECT

Aspen	Cottonwood	Hornbeam	Raspberry
Apple, American Crab	Curran	Horsechestnut	Rose, Cherokee
Barberry, Colorado	Elderberry	Juneberry	Sagebrush
Birch	Elder, Box	Locust, Black	Sassafras
Blackberry	Elm	Locust, Honey	Silverberry
Brambles	Gum, Black	Mesquite	Skunkbush
Buckbrush	Gum, Sweet	Mulberry	Shadbrush
Catalpa	Hackberry	Oaks	Succumb
Cedar, Salt	Hawthorn	Osage Orange	Sumac
Cherry, Chick	Haze	Poison Ivy	Sweetfern
Cherry, Wild	Herculesclub	Poison Oak	Wild Grape
Chestnut	Hickory	Poplar	Willow
Chinaberry	Honeysuckle	Tricky Ash	

If a knapsack sprayer is used dilute 1 to 2 pints of this material in 5 gallons of water and apply so that the plants are thoroughly wetted.

In the case of highly resistant woody plants such as Ash, Basswood, Beech, Red Maple, Mesquite, Persimmon, Rhododendron, Spruce, it is best to cut the plants to within a few inches of the ground and spray the stubble with a mixture of 1 part of this material diluted in 20 parts by volume of kerosene or diesel oil.

Do not graze dairy animals on treated areas within 7 days after application.

STUMP TREATMENT: To treat stumps of large trees or woody plants apply this material to freshly cut stumps diluted at the rate of 1 quart in 10 to 12 gallons of kerosene or diesel oil. This spray must be applied with a knapsack sprayer having a nozzle of medium orifice and care should be taken that both the sides and the tops of the stumps are thoroughly wetted. Should regrowth occur a second application may be necessary.

BRUSH CONTROL: This material can be used in Rangeland Clearance to control Sand Sagebrush, Scrub Oak, Mesquite and certain other noxious plants. Time of treatment is extremely important. It should be made when the plants are well leaved out and growing actively. 40-90 days before grazing.

PROCEDURE FOR RANGE LAND CLEARANCE

AIRPLANE APPLICATION: Mix 1 quart in 3 gallons of diesel oil and apply this volume per acre.

GROUND APPLICATION: Use 1 quart per acre in sufficient water or diesel oil depending upon equipment available, for thorough coverage.

NOTICE: The manufacturer makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on the manufacturer's label.

USDA Reg. No. 476-976

Made in U.S.A.

OOM670630MW

49

06 50