ACCEPTED
Jan 7 1983
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE, AND RODENTICIDE ACT FOR THE PRODUCT NAMED HEREON
EPA

Stauffer CHEMICAL

# TRITHION® 25-WP

## INSECTICIDE-ACARICIDE
## WETTABLE POWDER FOR USE IN WATER

| ACTIVE INGREDIENT: | |
|---|---|
| Carbophenothion, S-[p-chlorophenylthio)methyl] O,O diethyl phosphorodithioate | 25% |
| INERT INGREDIENTS: | 75% |
| | 100% |

## WARNING — Keep Out of Reach of Children

**MAY BE FATAL IF SWALLOWED.**

Do not get in eyes, on skin or clothing.
Avoid breathing dust and spray mist.
For commercial application, wear rubber gloves, dust respirator, protective clothing, rubber boots and goggles.
In case of contact, wash immediately with soap and water.
For eyes, flush with water for 15 minutes.
Wash hands, arms and face thoroughly with soap and water before eating or smoking.
Do not contaminate feed and food.
Wash contaminated clothing with soap and hot water before reuse.
ANTIDOTE — Exposure may cause cholinesterase inhibition. Call a physician immediately in all cases of suspected poisoning. Atropine is antidotal. 2-PAM may be antidotal when administered early and in conjunction with atropine.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

## USE PRECAUTIONS

**Read all precautions & directions before using.**

[illegible] Store in original container. Dispose of container [illegible]

## DIRECTIONS FOR USE

[illegible] can be made by aircraft or ground power equipment using [illegible] concentrate sprays. Only trained personnel should apply [illegible] Avoid spray drift in making applications. Do not apply with more than [illegible] recommended amount of this material on surfaces of [illegible] Fill the spray tank or premix in a bucket half filled with water [illegible] mixture through screen until nearly filled spray tank. Add [illegible] to fill tank. Keep agitator running during filling and spraying [illegible] Do not allow mixture to stand. Do not use in low volume [illegible] type spray equipment. If mixture does not mix readily, but forms [illegible] as an oily layer, do not use as injury to plants may result. Do not mix wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials has demonstrated them to be physically compatible and safe to plants.

## RECOMMENDATIONS

### VEGETABLE, FIELD AND VINE CROPS

Postbloom rates are [illegible] as pounds TRITHION 25-WP per acre. Unless otherwise specified below, apply the recommended dosage in sufficient [illegible] to provide thorough coverage according to spray equipment manufacturers' directions.

In California observe the following minimum water rate schedule:

| Crop | Gal. Water Per Acre Aircraft | Ground |
|---|---|---|
| Vegetable and Field Crops | 5 | 20 |
| [illegible] Fruits | 7 | 20 |

**ALFALFA (Fresh Forage and Hay):** Pea Aphid, Potato Leafhopper, Spotted Alfalfa Aphid, Two-Spotted Mite — Apply 2 to 4 pounds. Meadow Spittle Bug — Apply 4 pounds. Do not apply more than 1 application per cutting. Do not graze or cut hay within 28 days after application.

**BEANS, DRY (Shelled):** Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mite — Apply 2 pounds. Do not apply more than 2 applications per season. Do not graze, cut for forage or harvest beans within 21 days following the last application.

**BEANS, SNAP:** Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mite — Apply 2 pounds. Do not apply within 7 days of harvest. Do not make more than 2 applications per season. Do not graze or cut hay within 21 days after application.

**BEETS, GARDEN:** Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two-Spotted Mite — Use 1 to 2 pounds. Do not apply within 21 days of harvest if tops are to be used for food.

**CANTALOUPE, WATERMELONS:** Melon Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two Spotted Mite — Use 1 to 2 pounds. Do not apply within 5 days of harvest.

**CLOVER:** Pea Aphid, Potato Leafhopper, Two-Spotted Mite — Apply 2 to 4 pounds. Meadow Spittlebug — Apply 4 pounds. Do not make more than 1 application per season. Do not graze or cut hay within 28 days after application.

**CORN (Field and Sweet):** Grass Mite, Pacific Mite, Two Spotted Mite — Apply 4 pounds. Do not make more than 1 application per season. Do not graze or cut corn forage within 21 days after application.

**COTTON, Early Season:** Carmine Cinnabar, Mite, Two-Spotted Mite, Thrips — Use 1 to 2 pounds. **Mid and Late Season:** Use 1½ to 2 pounds. If mites persist, use high use 2 to 3 pounds and repeat in 5 to 10 days. **Cotton Leafworm** — Use 1 to 2 pounds. **Cotton Aphid, Leaf Perforator, Flea Hopper** — Use 4 pounds. Treated plants should not be grazed or fed to livestock.

**CUCUMBERS, SUMMER SQUASH:** Melon Aphid, Potato Leafhopper, Serpentine Leaf Miner, Two Spotted Mite — Use 2 to 4 pounds. Do not apply within 7 days of harvest.

**EGGPLANT, PEPPER, PIMIENTO, Mites (Tropical, Two-Spotted,** Potato Flea Beetle, Potato Leafhopper, Serpentine Leaf Miner — Use 2 to 4 pounds. Do not apply within 7 days of harvest.

(Continued on Back Panel)

LBS. NET

Made in U.S.A. By

**STAUFFER CHEMICAL COMPANY**

WESTPORT, CT 06880



## RECOMMENDATIONS (continued)

**GRAPES: Grape Leafhopper, Two-Spotted Mite** — [illegible] apply with a 28 days of harvest. Do not apply to [illegible] Do not graze, [illegible] Mustard and [illegible]

**ONIONS, [illegible] SEEDS, SETS OR TRANSPLANTS (Dry Bulb or Green):** [illegible] Furrow Treatment: Muck Soil [illegible] Mineral Soil — 3 to 4 pounds per acre. Apply at time of [illegible] transplanting by spraying in the planting furrow. Mix in 100 gallons of water and apply 3 gallons of suspension per 1,000 feet of furrow. Keep the mixture agitated. **Methocel Seed Coating Treatment:** For the production of pelleted seeds and [illegible] from Onion Magget. Treat dry Onion seeds with adequate amount of 5% Methocel, then tumble the Methocel treated seed in any commercial seed treater or barrel to which the seeds [illegible] thoroughly coated with TRITHION. Use [illegible] TRITHION 25-WP per 15 pounds of seed or 4 ounces per 1 pound of seed.

**PEAS: Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mite** — Use 2 to 4 pounds. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

**SEED CROPS (Alfalfa, Clover, Corn, Sorghum, Cucurbits, Melons, Sugar Beets and Vegetable crops listed on this label) Aphids (Bean, Pea, Spotted Alfalfa, Melon), Mites (Brown Wheat, Two-Spotted), Serpentine Leaf Miner** — Use 2 to 4 pounds. Do not graze or feed crop residue to livestock. Sugar Beet Seed Crops can be treated up to 14 days of harvest and crop residue can be fed to livestock.

**SORGHUM: Two-Spotted Mite** — Apply 4 pounds. Do not graze or cut forage within 21 days after application. Use 1 application per season. Under certain climatic conditions and on some varieties of grain sorghum, organophosphorous chemicals may cause [illegible] to the leaves.

**SPINACH: Bean Aphid, Serpentine Leaf Miner, Two-Spotted Mite** — [illegible] to 4 pounds. Do not apply within 21 days of harvest.

**STRAWBERRY: Strawberry Aphid, Two-Spotted Mite** — [illegible] 2 pounds. Do not apply within 3 days of harvest.

**SUGAR BEETS: Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two-Spotted Mite** — 2 to 4 pounds. Do not apply within 7 days of harvest.

**TOMATO: Mites, Tomato Russet, Tropical, Two-Spotted), Potato Flea Beetle, Potato Leafhopper, Potato Psyllid, Serpentine Leaf Miner** — Use 2 to 4 pounds. Do not apply within 7 days of harvest.

### FRUIT AND NUT CROPS

Unless otherwise specified, pesticide dosage rates are given as pounds TRITHION 25-WP and gallons spray of where known, per 100 gallons water for use in a thorough coverage spray when used [illegible] concentrate sprays should be applied in sufficient water to provide coverage according to equipment manufacturers' directions.

The maximum permissible pesticide rate per acre, [illegible] TRITHION 25-WP, is given in parentheses. [illegible] must not be exceeded whether using dilute or concentrate sprays [illegible] a minimum of 4 pounds TRITHION 25-WP per acre.

In California observe the following minimum water rate schedule:

| Crop | Gal. Water Per Acre Aircraft | Ground |
|---|---|---|
| Orchard [illegible] Nature Trees | [illegible] | [illegible] |
| Citrus | [illegible] | 100 |

**ALMONDS (California only): Summer Sprays: Mites (Clover, Brown Almond, European Red, Two spotted)** — Use 1 pound. If almond hulls are to be used for feed do not apply within 7 days of harvest. Do not make more than 1 application per season. Multiple applications at intervals [illegible] may be made. Hulls are not to be used for feed. 10 pounds. **Dormant Sprays: Overwintering Mite and Aphid Eggs, Scales (European Fruit Lecanium, Olive (Parlatoria), San Jose** — Use 1 to 2 pounds plus 1 to 2 gallons spray oil.

**APPLES, CRABAPPLE, PEAR, QUINCE: Summer: Aphids, Apple, Rosy, Apple Woolly Aphid, Codling Moth, Mites (Clover, Brown Almond, European Red, Two Spotted)** — Use 1 pound. Apply only after cull fruit evaluation. Do not apply within 30 days of harvest. 5 pounds. **Dormant or Delayed Dormant: Overwintering Aphid and Mite Eggs, Pear Psylla, San Jose Scale** — Use 1 to 2 pounds plus 1 to 2 gallons spray oil with the exception of [illegible] to the leave.

**APRICOTS: Summer: Mites, Clover, Brown Almond, European Red, Two Spotted** — Use 1 pound. Do not apply within 30 days of harvest. 8 pounds. **Dormant: Overwintering Mite Eggs, Scales (European Fruit Lecanium, San Jose** — Use 1 to 2 pounds plus 1 to 2 gallons spray oil.

**CHERRY: Summer: Mites (Cherry Rust, Clover, Brown Almond, European Red, Two Spotted,** — Use 1 pound. Do not apply within 30 days of harvest (8 pounds). **Dormant: Overwintering Aphid and Mite Eggs, Scales, European Fruit Lecanium, Olive (Parlatoria — San Jose)** — [illegible] pounds plus 1 to 2 gallons spray oil.

**CITRUS: Mites (Citrus Red (Purple), Citrus Rust, Six-Spotted, Texas Citrus, Two-Spotted)** — Apply 1 to 1½ pounds per 100 gallons water. [illegible] pounds. Application may be made up to day of harvest. [illegible] pounds is used, do not apply within 14 days of harvest. [illegible] do not use) more than 1 of any [illegible] applications of spray [illegible] If more than one [illegible] made, do not reapply within 30 days of the previous [illegible] treat on Scales (California Red, Citricola, Cottony [illegible] Purple, Citrus Red, Yellow, Olive Parlatoria, Latania, Florida Red, Citrus Black, Cottony Cushion, Yellow, Olive Parlatoria, European Fruit Lecanium, Citrus Mealybug, Pseudococcus [illegible] Apply 1 to 2 pounds plus 1 to 2 gallons spray oil.

CAUTION: [illegible]

**POST-HARVEST APPLICATION TO DECIDUOUS FRUIT AND NUT [illegible]**

## RECOMMENDATIONS—NON-FOOD CROPS

**FOREST AND CHRISTMAS TREE PLANTINGS: European Pine Shoot Moth [illegible] Tip Moths** — Use [illegible]

**FARMS, TURF: Chinch Bugs, Rhodes Grass Scale, [illegible] Sod Webworm, Yellowstriped Armyworm** — [illegible]

**ORNAMENTALS:** [illegible] Flowers and Tuberous Rooted Flowers [illegible] Chrysanthemum, Compositae, Cornflower, Cosmos, Chrysanthemum, Dahlia, Delphinium, Fuchsia, Garden Flowers, Gardenia, Geranium, Jasmine, [illegible] Roses, Shrubs, Trees, Vines, Weeds, Bulb or [illegible]

**PRUNE: Summer: Leaf Curling Plum Aphid, Leafhoppers, Mites (Clover, Brown Almond, European Red, Two Spotted)** — Use 1 pound. Do not apply within 30 days of harvest. 8 pounds. **Dormant: Overwintering Mite and Aphid Eggs, Scales (European Fruit Lecanium, San Jose)** — Use 1 to 2 pounds plus 1 to 2 gallons spray oil.

**WALNUT: Summer: Codling Moth, Mites (Clover, Brown Almond, European Red, Two Spotted Mite), Walnut Aphid, and for suppression of [illegible]** — [illegible]

For Control of [illegible] Whitefly, Pine, Rose, [illegible] Mites, Thrips, Leafhoppers, Aphids, Mealybugs, Scales, Leafminers, Sawflies, Caterpillars, Beetles, Weevils, Borers, Midges, Gall Insects, Mites, Spiders, Ticks, Lacebugs, Psyllids, Florida Red Scale, Citrus Mites, Thrips, Whitefly, Citrus Blackfly, Purple Scale, Citrus Thrips, Yellow Scale, Citrus Mealybug, Glover Scale, Citrus Red Mite, Texas Citrus Mite, Citrus Rust Mite, Six-Spotted Mite, Two-Spotted Mite, Citrus Bud Mite, Cyclamen Mite, Cottony Cushion, Yellow, Olive Parlatoria, European Fruit Lecanium, Citrus Mealybug, Pseudococcus, Citrophilus, Latania, Purple Scale, Citrus Red Scale, Yellow Scale, Citrus Blackfly, Citrus Thrips, Whitefly, Glover Scale, Citrus Red Mite, Texas Citrus Mite, Citrus Rust Mite, Six-Spotted Mite, Two-Spotted Mite, Citrus Bud Mite, Cyclamen Mite, Broad Mite, Privet Mite, Boxwood Mite, Pine Needle Scale, Juniper Scale, Oystershell Scale, Lecanium Scale, Euonymus Scale, Tuliptree Scale, Obscure Scale, Calico Scale, Fletcher Scale, Cooley Spruce Gall Aphid, Eastern Spruce Gall Aphid, Balsam Twig Aphid, Spruce Spider Mite, Juniper Spider Mite, Arborvitae Leafminer, Pine Needle Miner, Birch Leafminer, Elm Leafminer, Spruce Needle Miner, Pine Needle Miner, Pine Tip Moth, Eastern Pine Shoot Moth, Zimmerman Pine Moth, Pine Tube Moth, Sawflies, Pine Sawfly, Larch Sawfly, Spruce Sawfly, Birch Leaf Miner, Elm Leaf Miner