

# CAL-META<sup>®</sup> SNAIL PELLETS

AGRICULTURAL POISON BAIT

7022  
Sep 27, 1971  
UNDER THE  
FEDERAL  
ED. LINE ER 476-1092

**Active Ingredients:**

Tricalcium Arsenate .....	6.75%
Metaaldehyde .....	1.50%
Inert Ingredients .....	91.75%
	<b>100.00%</b>

Arsenic, expressed as metallic ..... 2.54%  
Arsenic in water soluble form, expressed as metallic, not more than ..... 2.54%

## WARNING

Keep Out of Reach of Children

### HARMFUL IF SWALLOWED

AVOID CONTACT WITH SKIN, EYES OR CLOTHING.  
DO NOT CONTAMINATE FOOD OR FEED.  
FIELD WORKERS SHOULD USE PROTECTIVE GLOVES WHEN HANDLING  
AND BROADCASTING CAL-META SNAIL PELLETS.  
WASH WITH SOAP AND WATER AFTER HANDLING.  
THIS BAIT MAY BE ATTRACTIVE TO DOGS. KEEP DOGS OUT OF  
TREATED AREAS.

® CAL-META is a registered trade-mark of Stauffer Chemical Company.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-1092  
ARC-710803-6

Made in U. S. A.  
Stauffer Chemical Company  
New York

50 LB. NET

## RECOMMENDATIONS

CAL-META SNAIL PELLETS is a special formulation which employs apple chop as the base carrier for the main toxicants, calcium arsenate and metaaldehyde. After careful blending and mixing the material is processed through special extruding equipment to produce a pelleted product of uniform composition and consistency.

CAL-META SNAIL PELLETS are recommended for control of SLUGS, SNAILS and CUTWORMS in asparagus, beans, blackeyed peas, broccoli, Brussels sprouts, cabbage, carrots, cauliflower, celery, collards, corn, cucumbers, eggplants, kale, kohlrabi, melons, peppers, pumpkins, rutabagas, spinach, squash (summer and winter), strawberries (do not apply after fruit starts to form), tomatoes, turnips, and ornamentals.

**DIRECTIONS** Apply 30-50 lb. per acre.

CUTWORMS, SNAILS, SLUGS: Application can be made by hand or power equipment. Treat infested areas. Repeat when necessary.

SLUGS AND SNAILS: For hand application, broadcast liberally under trees, around irrigation standpipes, buildings and other places where slugs and snails concentrate. Repeat application following irrigation. Apply in evening.

**CAUTION—DO NOT APPLY DIRECTLY TO FOLIAGE OR AGAINST THE BASE OF PLANTS.** Burning may result from bait becoming lodged in plant crevices. Apply only to soil surface around plants. Do not apply to foliage or other edible parts.

TOXIC TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. BIRDS FEEDING ON TREATED AREAS MAY BE KILLED.

**IMPORTANT: Protect life and domestic animals. Burn this bag when empty—Bury the ashes.**