

# CROWN® 90 WETTABLE SULFUR

Crown 90 Wettable Sulfur is formulated for use in water as a spray on fruit, vegetable and ornamental crops for the control of certain fungus diseases and mites. It may be combined in the spray tank with most of the commonly used fungicides and insecticides.

## DIRECTIONS FOR USE

Pour the specified amount of this product on surface of water in nearly filled spray tank or pre-mix in a bucket before pouring into spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine with emulsifiable liquids unless previous use of the mixture has proven physically compatible and safe to plants.

## RECOMMENDATIONS

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

All dosages given on this label are for dilute sprays (IX concentrations). For use in concentrate sprays, increase the dosage of Crown 90 Wettable Sulfur proportionately. Spray often enough to keep new growth covered.

**APPLE, PEAR: Powdery Mildew, Scab-** Use 6 to 8 lb. per 100 gal. of water in the pre-bloom, bloom and petal fall sprays. Use 4 to 6 lb. in the cover sprays.

**CHERRY: Brown Rot, Leaf Spot-** Use 6 to 8 lb. per 100 gal. of water in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

**CITRUS: Rust Mite-** Use 5 lb. per 100 gal. of water. Begin applications at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russetting of fruit and greasy spot on leaves. Scale Crawlers -- Add 5 to 8 lb. to each 100 gal. of tank strength lime sulfur solution.

**PEACH: Brown Rot, Scab-** Use 6 to 8 lb. per 100 gal. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**PLUM, PRUNE: Brown Rot, Leaf Spot-** Use 6 to 8 lb. per 100 gal. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**BEANS, PEAS: Powdery Mildew, Rust-** Use 16 lb. per acre in sufficient water for thorough coverage. Repeat applications in 7 to 14 day intervals as necessary.

**ROSES, PERENNIALS, SHRUBS: Black Spot, Powdery Mildew, Rust-** Use 6 to 8 lb. per 100 gal. of water (3 to 4 tablespoons per 1 gal. of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

**IMPORTANT:** Use the higher dosage during periods of frequent and heavy rainfall. Spray often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Crown 90 always add the same amount of Spray Lime, or other safener, as would be used when applying Lead Arsenate without sulfur. When using Zinc Sulfate and Spray Lime as a safener on peaches, always add Zinc Sulfate to the spray tank first, followed by Spray Lime, Lead Arsenate and sulfur in order given.

**CAUTION:** Do not use sulfur in combination with oil or within two weeks of an oil spray. Do not use sulfur on apples or other sensitive crops when shade temperatures exceed, or are likely to exceed, 90° F. Do not apply or allow sulfur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tung. To do so may result in severe fruit and leaf injury.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label. 2-ARC-730419-0

**DO NOT RE-USE CONTAINER, DESTROY WHEN EMPTY.**



# CROWN 90

WETTABLE

# SULFUR

FOR SPRAYING FRUITS,  
VEGETABLES, ORNAMENTALS

ACTIVE INGREDIENT: SULFUR . . . . . 90%

INERT INGREDIENTS . . . . . 10%

CROWN is a registered trademark of Stauffer Chemical Company. **100%**

FINENESS: not less than 93% passing 325 mesh

**CAUTION**

EPA Reg. No. 476-125 AA

KEEP OUT OF REACH OF CHILDREN

Made in U.S.A. by

## STAUFFER CHEMICAL COMPANY

### WESTPORT, CT 06880



CROWN 90  
SULFUR

### 30 LB.NET

ACCEPTED  
AUG 31  
DILLON  
FOR  
28243

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information as the timing and number of applications needed will vary with local conditions.

All dosages given on this label are for dilute sprays (IX concentrations). For use in concentrate sprays, increase the dosage of Crown 90 Wettable Sulfur proportionately. Spray often enough to keep new growth covered.

**APPLE, PEAR: Powdery Mildew, Scab-** Use 6 to 8 lb. per 100 gal. of water in the pre-bloom, bloom and petal fall sprays. Use 4 to 6 lb. in the cover sprays.

**CHERRY: Brown Rot, Leaf Spot-** Use 6 to 8 lb. per 100 gal. of water in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

**CITRUS: Rust Mite-** Use 5 lb. per 100 gal. of water. Begin applications at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russeting of fruit and greasy spot on leaves. Scale Crawlers - Add 5 to 8 lb. to each 100 gal. of tank strength lime sulfur solution.

**PEACH: Brown Rot, Scab-** Use 6 to 8 lb. per 100 gal. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**PLUM, PRUNE: Brown Rot, Leaf Spot-** Use 6 to 8 lb. per 100 gal. of water in pink, bloom, shuck, cover and pre-harvest sprays.

**BEANS, PEAS: Powdery Mildew, Rust-** Use 16 lb. per acre in sufficient water for thorough coverage. Repeat applications in 7 to 14 day intervals as necessary.

**ROSES, PERENNIALS, SHRUBS: Black Spot, Powdery Mildew, Rust-** Use 6 to 8 lb. per 100 gal. of water (3 to 4 tablespoons per 1 gal. of water). Make first application prior to first signs of disease and maintain preventive schedule throughout the season, repeating applications every 7 to 14 days. Keep new growth protected.

**IMPORTANT:** Use the higher dosage during periods of frequent and heavy rainfall. Spray often enough to keep susceptible growth covered. This may require weekly or bi-weekly applications during periods of rapid growth and frequent rainfall. When using Lead Arsenate with Crown 90 always add the same amount of Spray Lime, or other safener, as would be used when applying Lead Arsenate without sulfur. When using Zinc Sulfate and Spray Lime as a safener on peaches, always add Zinc Sulfate to the spray tank first, followed by Spray Lime, Lead Arsenate and sulfur in order given.

**CAUTION:** Do not use sulfur in combination with oil or within two weeks of an oil spray. Do not use sulfur on apples or other sensitive crops when shade temperatures exceed, or are likely to exceed, 90° F. Do not apply or allow sulfur to drift to Cucurbits (cucumber, cantaloupe, melon, squash) or tung. To do so may result in severe fruit and leaf injury.

**NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.** 2-ARC-730419-0

**DO NOT RE-USE CONTAINER, DESTROY WHEN EMPTY.**

FILMED - 1978

Rehab Group

(703) 521-7000