

# ZINEB 5 DUST


## AN AGRICULTURAL FUNGICIDE

### ACTIVE INGREDIENT:

Zineb (Zinc Ethylene Bisdithiocarbamate) . . . . . 5%

### INERT INGREDIENTS: . . . . . 95%

Zinc expressed as metallic . . . . . 100%



**CAUTION** Keep Out of Reach of Children

- DO NOT BREATHE DUST.
- MAY CAUSE IRRITATION TO NOSE AND THROAT.

### USE PRECAUTIONS

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified.

Consult State Agricultural Extension Service or State Agricultural Experiment Stations for additional information, as the timing, number, and rate or applications needed will vary with local conditions.

### DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infection and repeat at 5 to 10 day intervals, depending on severity of infection, but observe use limits given for specific crops. Apply at the rate of 30 to 40 lb. per acre unless otherwise specified.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

### RECOMMENDATIONS

- BEANS** --- Anthracnose, Downy Mildew, Rust. Do not apply within 7 days of harvest on lima, snap or string beans.
- CABBAGE** --- Downy Mildew, Black or Leaf Spot. Do not apply within 7 days of harvest.
- GARROTS** --- Early Blight, Late Blight. Do not apply within 7 days of harvest if created tops are to be used as food or feed.
- VEGETABLES** --- Early and Late Blight. Remove residues by washing, stripping, trimming. *Early Blight*
- (Cucumbers, Melons, Pumpkins, Squash)** --- Angular Leaf Spot, Anthracnose, Downy Mildew, Scab.
- EGGPLANT and PEPPERS** --- Alternaria, Anthracnose, Cercospora, Phomopsis.
- GRAPES** --- Black Rot, Bunch Rot, Downy Mildew. Do not apply within 7 days of harvest.
- LETTUCE** --- Downy Mildew. Do not apply on leaf lettuce within 7 days of harvest and on head lettuce within 5 days of harvest.
- ONIONS** --- Botrytis Leaf Blight, Downy Mildew, Purple Blotch. Do not apply on green onions within 7 days of harvest.
- PEANUTS:** Cercospora Leaf Spot. Apply 25 lb. per acre. Do not use treated peanut hay for feed.
- SPINACH** --- White Rust, Blue Mold. Do not apply within 7 days of harvest.
- STRAWBERRIES** --- Leaf Spot. Do not apply within 7 days of harvest.
- TOMATOES** --- Anthracnose, Early and Late Blight, Gray Leaf Spot (Stemphylium), Septoria Leaf Spot. *Early Blight*
- AZALEAS** --- Petal Blight. (Make applications to the flowers, soil, and litter around the plant beginning with the bud swelling stage and continue through the flowering season).
- ROSES** --- Black Spot.