

300
17
138127
1

Stauffer Chemical Company
1200 S. 47th Street
Richmond, CA 94804

OCT 3 1984

Attention: Ralph L. Riggs
Pesticide Registrations

Gentlemen:

Subject: Folpet (Phaltan) 75-WP
EPA Registration No. 476-1294
Your Letter Dated August 15, 1984

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted at the next printing.

1. Add the environmental hazards phrase, "Do not apply directly to water ~~or wetlands~~," after the fish toxicity statement.
2. Change "Caution" to "Note to User" in the Directions for Use.

A stamped copy of the label is enclosed for your records.

Sincerely yours,



Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

RD/FHB:Jacobey:DCR-32907:0165L:efs:Raven:479-2013:03/5/84:Del.9/20/84

CONCURRENCES

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

(No type on this label will be less than 6 point)

(Stauffer Shield)

FOLPET (PHALTAN*) 75-WP

Agricultural Fungicide For Plant Disease Control
Wettable Powder For Use In Water

ACTIVE INGREDIENT:

FOLPET: N-(trichloromethylthio)phthalimide 75%

INERT INGREDIENTS: 25%
100%

Contains 75 pounds of active ingredient per 100 pounds.

*PHALTAN is a registered trademark for phthalimide fungicide

KEEP OUT OF REACH OF CHILDREN

REGISTERED
WITH COMMENTS
EPA Letter Dated:

C A U T I O N

OCT 3 1984

STATEMENT OF PRACTICAL TREATMENT

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended,
registered

476-1394

First Aid

Immediately start the procedures given below and contact a Poison Center, a physician or the nearest hospital for further assistance or call Collect, Day or Night, (203) 226-6602 Stauffer Chemical Company or (800) 424-9300 Pesticide Team Safety Network.

If swallowed, immediately give several glasses of water and induce vomiting by gagging the victim with a finger placed at the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious or convulsing, do not induce vomiting or give anything by mouth.

For skin contact, flush all affected areas with plenty of water for several minutes. Remove and clean contaminated clothing and shoes. Seek medical attention if irritation occurs.

For eye contact, immediately flush eyes with large amounts of running water for at least 15 minutes. Hold eyelids apart during the flushing to ensure rinsing entire surface of the eye and lids with water. Get medical attention immediately.

If inhaled, remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

_____ LBS. Net _____ KG

EPA Reg. No. 476-1294
EPA Est. 476-

RS-840727

This product is sold only for uses stated on its label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country: United States.

Made in U.S.A. by
STAUFFER CHEMICAL COMPANY
Westport, Ct. 06881-0850

CELERY: Cercospora leaf spot, early and late blights. Use 1-1/3 to 2-2/3 pounds per acre in sufficient water for thorough coverage. Make first application after plants are set in field and established. Repeat at 7 day intervals. Do not apply within 7 days of harvest. (2-2/3 pounds)

CHERRIES (RED TART): Brown rot. Use 2/3 to 1-1/3 pounds. Apply in delayed dormant, popcorn, and bloom stages and at 7 to 14 day intervals; thereafter, as necessary. Leaf spot. Use 2/3 to 1-1/3 pounds. Apply in cover period at 7 to 14 day intervals as necessary and repeat once or twice as post-harvest foliage applications. Do not use on sweet cherries as foliage injury may occur. (8 pounds)

CITRUS (GRAPEFRUIT, KUMQUATS, LEMONS, LIMES, ORANGES, TANGELOES, TANGERINES): Greasy spot. Use 1-1/3 pounds plus sticker. Apply in August and September, 3 to 10 weeks after first growth. Melanose. Use 1-1/3 to 2-2/3 pounds. Make first application 1 week after bloom and complete as soon as possible. Scab. Use 1-1/3 pounds. Apply in dormant and 2/3 petal fall stages, 2 weeks later, and on fall flush in August and September. (26-2/3 pounds)

CRANBERRIES: Storage rots. Use 6 pounds per acre, in sufficient water for thorough coverage. Apply during bloom and repeat at 10 to 14 day intervals. Do not apply within 30 days of harvest. (12 pounds)

CUCUMBERS: Anthracnose, downy mildew, fruit rots (pythium and rhizoctonia), powdery mildew. Use 1-1/3 to 2-2/3 pounds. Apply 200 gallons per acre. Make first application when first true leaves appear. Repeat at weekly intervals. (5-1/3 pounds)

CURRENTS, GOOSEBERRIES: Leaf spot. Use 2-2/3 pounds. Apply 150 gallons per acre. Make first application before infection or before bloom. Repeat at 7 to 10 day intervals in the post-bloom period. (4 pounds)

GRAPES: Black rot, downy mildew. Use 1-1/3 pounds. Apply 250 to 300 gallons per acre. Make first application just before bloom. Repeat just after bloom and at 7 to 10 day intervals for 1 to 3 more applications. Dead-arm. Use 1-1/3 pounds. In eastern states, begin when new growth is 1 to 2 inches long. Repeat when 8 to 10 inches long. In western states, begin at bud break. Repeat 2 weeks later or when new growth is 4 to 8 inches long. Powdery mildew. Use 1-1/3 pounds. Begin 2 to 3 weeks after bloom. Repeat at 2 week intervals. NOTE: During periods of extended hot, dry weather foliage injury may occur. (4 pounds)

LETTUCE: Downy mildew. Use 1-1/3 pounds. Apply up to 200 gallons per acre. Make first application when plants are established in field. Repeat at 7 day intervals until harvest. (2-2/3 pounds)

ONIONS, GARLIC, LEEKS, SHALLOTS: Downy mildew, purple blotch. Use 2-2/3 pounds. Apply 200 gallons per acre. Make first application before infection. Repeat at 7 day intervals until tops begin to dry before harvest. (5-1/3 pounds)

STRAWBERRIES: Fruit rot, leaf spot. Use 1-1/3 pounds. Apply up to 200 gallons per acre. Make first application before bloom and repeat at 7 day intervals until after harvest. (2-1/3 pounds)

TOMATOES: Anthracnose. Use 2-1/3 pounds. Apply up to 200 gallons per acre. Make first application during bloom and repeat at 7 to 10 day intervals through harvest. (5-1/3 pounds)

ORNAMENTALS

Unless otherwise specified dosage rates are in terms of FOLPET (PHALTAN) 75-WP per 100 gallons of water. Thorough coverage is required.

ASTERS, PHLOX: Powdery mildew. Use 1-1/3 pounds. (1-1/3 tablespoonfuls per 1 gallon water.) Make first application before disease appears and repeat at 7 to 10 day intervals. Increase frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

AZALEAS (CUTTINGS): Damping-off. Use 1 to 1-1/3 pounds. Soak cuttings for 15 to 30 minutes before placing in rooting media.

CARNATIONS: Alternaria leaf spot, rust. Use 2/3 to 1-2/3 pounds. (2/3 to 1-2/3 tablespoonfuls per 1 gallon water.) Make first application 2 weeks after planting and repeat at 7 to 10 day intervals as necessary.

CHRYSANTHEMUMS: Powdery mildew, septoria leaf spot. Use 1-2/3 pounds. (1-2/3 tablespoonfuls per 1 gallon water.) Make first application immediately after planting and repeat at 7 to 10 intervals. Increase frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

GLADIOLUS (CORNS): Corn rot, damping-off. Use 10 teaspoonfuls per 1 gallon water. Apply to corns as a pre-planting soak for 20 to 30 minutes. Keep mixture stirred. Drain and plant immediately.

GRASS (BLUEGRASS IN NON-PASTURED SITES): Melting out (helminthosporium). Use 4 ounces in 5 gallons water per 1000 square feet. Make first application before disease is expected to appear and repeat at 7 day intervals as necessary. Do not graze treated areas. Do not feed clippings to livestock.

IRIS: Didymellina leaf spot. Use 1-2/3 pounds. (1-2/3 table-spoonfuls per 1 gallon water.) Make first application when plants emerge and repeat at 7 to 10 day intervals.

MARIGOLDS, ZINNIAS: Alternaria leaf spot. Use 1-2/3 pounds. (1-2/3 tablespoonfuls per 1 gallon water.) Make first application when plants emerge and repeat at 3 to 7 day intervals. Use higher frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

POINSETTIAS (GREENHOUSE): Pythium, root rot. Use 1-2/3 pounds plus a spreader and use 1 pint of suspension per 8 inch pan. Make application at time of panning and repeat 10 days later.

ROSES: Black spot, powdery mildew. Use 2/3 to 1-2/3 pounds. (2/3 to 1-2/3 tablespoonfuls per 1 gallon water.) Make first application when first leaves unfold and repeat at 7 to 10 day intervals throughout the season. Do not use wetting agents on roses as injury may result.

SNOWDRAGONS: Anthracnose, powdery mildew, rust. Use 1-2/3 pounds. (1-2/3 tablespoonfuls per 1 gallon water.) Make first application when plants emerge and repeat at 3 to 7 day intervals. Use higher frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in cool, dry place. Protect from excessive heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE - READ CAREFULLY

CONDITIONS OF SALE

Stauffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION

Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties; THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assume all risks associated with use or application of this product contrary to label instruction or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY

In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

GRASS (BLUEGRASS IN NON-PASTURED SITES): Melting out (helminthosporium). Use 4 ounces in 5 gallons water per 1000 square feet. Make first application before disease is expected to appear and repeat at 7 day intervals as necessary. Do not graze treated areas. Do not feed clippings to livestock.

IRIS: Didymellina leaf spot. Use 1-2/3 pounds. (1-2/3 tablespoonfuls per 1 gallon water.) Make first application when plants emerge and repeat at 7 to 10 day intervals.

MARIGOLDS, ZINNIAS: Alternaria leaf spot. Use 1-2/3 pounds. (1-2/3 tablespoonfuls per 1 gallon water.) Make first application when plants emerge and repeat at 3 to 7 day intervals. Use higher frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

POINSETTIAS (GREENHOUSE): Pythium, root rot. Use 1-2/3 pounds plus a spreader and use 1 pint of suspension per 8 inch pan. Make application at time of panning and repeat 10 days later.

ROSES: Black spot, powdery mildew. Use 2/3 to 1-2/3 pounds. (2/3 to 1-2/3 tablespoonfuls per 1 gallon water.) Make first application when first leaves unfold and repeat at 7 to 10 day intervals throughout the season. Do not use wetting agents on roses as injury may result.

SNAPDRAGONS: Anthracoze, powdery mildew, rust. Use 1-2/3 pounds. (1-2/3 tablespoonfuls per 1 gallon water.) Make first application when plants emerge and repeat at 3 to 7 day intervals. Use higher frequency during periods of rapid new growth and severe infection caused by rains or high humidity.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in cool, dry place. Protect from excessive heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE - READ CAREFULLY**CONDITIONS OF SALE**

Stauffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be varied by written agreement of a duly authorized representative of Stauffer.

WARRANTY LIMITATION

Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties; THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

INHERENT RISKS

The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assume all risks associated with use or application of this product contrary to label instruction or resulting from extraordinary weather conditions.

LIMITATION OF LIABILITY

In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.