



TRITHION™ 10-G

GRANULES — SOIL INSECTICIDE

476-1346
ACCEPTED
Mar 1 1973

ACTIVE INGREDIENT:

Carbophenothion S-[(p-chlorophenylthio)methyl]
O,O-diethyl phosphorodithioate

10%

INERT INGREDIENTS

90%

100%

U.S. Patent No. 2,797,224

OBSERVE THE FOLLOWING USE PRECAUTIONS

Read all precautions and directions before using.
Use only for claims listed and only as specified on this label.
In order that pesticides used for food and forage crops will not exceed tolerances established by the Environmental Protection Agency, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas used by unprotected humans or domestic animals or onto a fumigant, fertilizer or pesticide. **The grower is responsible** for residue on his crops and for damage caused by drift from his property to that of others.

FISH AND WILDLIFE CAUTIONS: Toxic to fish and wildlife. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Birds feeding in treated areas may be killed.

RECOMMENDATIONS AND DIRECTIONS FOR USE

ONIONS: From seeds, sets or transplants (dry bulb or green) — Onion Maggot Control: Furrow Treatment— Apply in the planting furrow at time of seeding or transplanting using a combination drill or suitable granular application equipment.

Muck Soil—Apply 2 1/2 pounds per acre for planting in 18-inch rows (21 ounces per 100 ft. row) or 10 pounds per acre for planting in 18-inch rows (11 pounds per 100 ft. row). Do not exceed the recommended dosage rate.

Mineral Soil—Apply 10 to 15 pounds per acre for planting in 18-inch rows (11 to 15 ounces per 100 ft. row).

SUGAR BEETS: Sugar Beet Root Maggot (Tetanops Myopaeformis)—Apply in the reduction of Sugar Beet Root Maggot, apply 10 pounds TRITHION 10-G per acre in the drill row at seeding time just above the seed. Use a combination drill or suitable granular application equipment.

LAWN AND TURF: Chinch Bugs—Apply 3 1/2 pounds of TRITHION 10-G per 1,000 square feet (40 1/2 lbs. of lawn maintenance fertilizer spreaders when using the number used for maintenance seed with granular fertilizer). Distribute the material uniformly, after application, water with water sprinkler thoroughly. Do not permit the water to dry before the material has been distributed.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED.

- Avoid breathing dust, contact with skin and eyes.
- Wear dust respirator and goggles.
- Wash hands, arms and face thoroughly with soap and water before eating or smoking.
- Do not contaminate feed and food.
- Wash all contaminated clothing with soap and hot water before reuse.

ANTIDOTE—Exposure may cause cholinesterase inhibition. Call a physician immediately in all cases of suspected poisoning. Atropine is antidote. 2-PAM is also antidotal when administered early, and for combination with atropine.

BURY WASTE AND EMPTY CONTAINER IN AN ISOLATED AREA LOCATED AWAY FROM WATER SUPPLIES

POUNDS NET

EPA Reg. No. 476-1346

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880