ACCEPTED

UNDER THE FEDERAL INSECTICIDE ACT
FOR REGISTERED USES
SUBJECT TO ALL WARNINGS

USE PRECAUTIONS

Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed tolerances established by the Environmental Protection Agency, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or livestock animals or onto adjoining food, fiber or pasture crops.

The grower is responsible for residues on his crop as well as for damages caused by drift from his property to that of others.

Consult State Agricultural Experiment Station or State Agricultural Extension Service for additional information as the timing and number of applications will vary with local conditions.

Do not apply sprays to trees where soil moisture is low or has recently been low.

Do not apply sprays when temperatures are in excess of 90° F. or in periods of low humidity, caused by hot winds.

Do not combine sprays with sulfur sprays or apply within 30 days before or after sulfur.

FISH AND WILDLIFE CAUTIONS: This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Shrimp and crab may be killed at low concentrations recommended on this label. Do not apply where they are important resources. Apply this product as specified on the label.

To protect [illegible] do not apply when bees are actively foraging in the area to be treated.

DIRECTIONS FOR USE

Applications may be made by aircraft or ground equipment using sufficient volume of concentrate sprays. Only trained persons using approved protective equipment should make applications. Do not apply with hand equipment. Fill specified amount of this product into partially filled spray tank. Add balance of water to fill tank. Keep agitator running during filling and spraying operations. Do not allow material to stand. If material does not mix readily or tends to separate as an oily layer, do not use as many to plants may result. Do not combine with emulsifiable type pesticides or wettable powders unless previous use of the mixture has proven physically compatible and safe to plants.

RECOMMENDATIONS

VEGETABLE, FIELD AND VINE CROPS

Each side rate given as pints TRITHION 4-E per acre (unless otherwise specified) should be applied in the recommended dosage in enough water to provide adequate coverage. Consult spray equipment manufacturers' directions for gallons thereof. Use the [illegible] **minimum** water rate schedule.

CROP	Gal. Water Per Acre Aircraft	Ground
[illegible]	5	[illegible]
Small Fruits	7	20

ALFALFA (Fresh Forage and Hay), CLOVER: Pea Aphid, Potato Leafhopper, Spotted Alfalfa Aphid, Two Spotted Mite — Apply 1 to 2 pints. Meadow Spittlebug — Apply 2 pints. Do not apply more than one application per cutting. Do not graze or cut for hay within 28 days after application.

BEANS, LIMA: Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpentine Leaf Miner, Two Spotted Mite — Apply 1 to 1½ pints. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

BEANS, SNAP: Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mite — Apply 1 pint. Do not apply within 7 days of harvest. Do not make more than two applications per season. Do not graze or cut for hay within 21 days after application.

BEANS, DRY: Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mite — Apply 1 pint per acre. Do not make more than two applications per season. Do not graze, cut for forage or harvest beans within 21 days following the last application.

BEETS, GARDEN: Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two-Spotted Mite — Use ½ to 1 pint. Do not apply within 21 days of harvest if tops are to be used as feed or food.

CANTALOUPE, WATERMELONS: Melon Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two-Spotted Mite — Use 1 to 2 pints. Do not apply within 5 days of harvest.

CORN (Field and Sweet): Grass Mite, Pacific Mite, Two-Spotted Mite — Apply 2 pints. Do not make more than one application per season. Do not graze or cut for forage within 21 days after application.

COTTON: Early Season — Carmine Cinnabar Mite, Two Spotted Mite, Tropical Mite — Use ½ to 1 pint. **Mid and Late Season** — Use ½ to 1 pint. If mite populations are high use 1 to 1½ pints and repeat in 5 to 10 days. Cotton Leafworm — Use ½ to 1 pint. Cotton Aphid, Leaf Perforator, Fleahopper — Use 2 pints per acre. Treated plants should not be grazed or fed to livestock.

CUCUMBERS, SUMMER SQUASH: Melon Aphid, Potato Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mite — Use 1 to 2 pints. Do not apply within 7 days of harvest.

EGGPLANT, PEPPER, PIMENTO: Mites, Tropical, Two Spotted, Potato Flea Beetle, Potato Leafhopper, Serpentine Leaf Miner — Use 1 to 2 pints. Do not apply within 7 days of harvest.

GRAPES: Grape Leafhopper, Two Spotted Mite — 1 pint. Do not apply within 30 days of harvest. Do not apply to sensitive varieties such as Charbono, Gamay, Golden Muscat, Muscat and Hamburg.

ONION: From Seeds, Sets or Transplants (for Dry Bulb or Green): Onion Maggot control — [illegible] treatment. **Muck Soils:** 3 to 4 pints per acre. **Mineral Soil:** 2 to 3 pints per acre. Apply at time of seeding or transplanting by spraying in the planting furrow. Mix in 100 gallons of water and apply gallons of mixture per [illegible] of furrow. Keep the mixture agitated.

PEAS: Pea Aphid, Serpentine Leaf Miner, Two Spotted Mite — Use ½ to 1 pints. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

CROPS [illegible] Sugarbeets and Vegetable Crops listed on this label): [illegible] Spotted Mite, Mites (Brown Wheat, [illegible]) — Use 1 to [illegible] pints. Do not graze or feed crop residue to livestock. Sugarbeet seed crops can be treated up to 14 days of harvest and crop residue can be fed to livestock.

SORGHUM: Green Bug, Greenbug, Pea Aphid, Sorghum Midge, Two Spotted Mites, [illegible] Mite Species — 1 pint. Limit applications to two per season. Do not graze or cut for forage within 21 days. Under certain climatic conditions, and on some varieties of grain sorghum, organophosphate chemicals may cause a slight discoloration to the leaves.

SOYBEANS: Bean Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpentine Leaf Miner, Two Spotted Mite. Apply 1 to 1½ pints. Do not apply within 7 days of harvest. Do not feed treated forage to livestock.

SPINACH: Bean Aphid, Serpentine Leaf Miner, Two Spotted Mite — Use ½ to 1 pint. Do not apply within 21 days of harvest.

STRAWBERRY: Strawberry Aphid, Two Spotted Mite — Use ½ pint. Do not apply within 3 days of harvest.

SUGARBEETS: Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two Spotted Mite — Use 1 to 2 pints. Do not apply within 14 days of harvest.

TOMATO: [illegible] Tropical, Two Spotted, Potato Flea Beetle, Potato [illegible], Potato Psyllid, Serpentine Leaf Miner — Use 1 to [illegible] pints. Do not apply within 7 days of harvest.

Stauffer

TRITHION® 4-E

EMULSIFIABLE LIQUID INSECTICIDE-ACARICIDE

CONTAINS FOUR POUNDS TRITHION PER GALLON

ACTIVE INGREDIENTS:	
Carbophenothion S-[(p-chlorophenylthio)methyl] O,O-diethyl phosphorodithioate	45.9%
Xylene Range Aromatic Hydrocarbon Solvent	45.5%
INERT INGREDIENTS:	8.6%
	100.0%

Rinse spray equipment and containers. Discharge liquid waste in pit in isolated area located away from water supplies.

PERFORATE EMPTY METAL CONTAINERS. CRUSH AND BURY GLASS.

U.S. Patent No. 2,749,223

WARNING

Keep Out of Reach of Children

MAY BE FATAL IF SWALLOWED.

Do not get in eyes, on skin or on clothing. Avoid [illegible] application. Wear rubber gloves, protective clothing, rubber overshoes and goggles. In case of contact, wash immediately with soap and water. In case of eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Do not contaminate food. Wash contaminated clothing before use and hot water before re-use.

ANTIDOTE — Exposure may cause [illegible] symptoms. [illegible] immediate medical attention. In cases of suspected poisoning, Atropine sulfate [illegible] should be administered orally and in conjunction with atropine.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Protect from temperatures below 0° F. Liquid phase may separate at lower temperatures. Warm or store at room temperature and mix thoroughly before use.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

COMBUSTIBLE! Do Not Use or Store Near Heat or Open Flame.

LOT NO. GALLONS NET

EPA REG. NO. 476-1376-AA

Made In U.S.A. By [illegible]

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

FRUIT AND NUT CROPS

Unless otherwise specified, for foliage dormant sprays where shown, TRITHION 4-E and [illegible] sprays or where shown, use a thorough coverage spray. When used as directed, sprays should be applied in sufficient water to equipment manufacturers' directions. The maximum permissible pesticide rate per acre is given in parenthesis [illegible] after each crop, whether using [illegible] amount of TRITHION 4-E per acre. [illegible] rate schedule.

CROP	
[illegible] Mites, [illegible]	
Almonds	California only: **Pre-Bloom or Delayed Dormant:** [illegible] pint. **Summer:** Mites, [illegible] Use ½ pint plus up to 2 gallons of oil. Used for feeding, do not apply within 7 days. One application per season. Multiple applications may be made if buds are not fully developed. Overwintering Mite and Aphid Eggs, Scale Crawlers — Use ½ to 1 pint.
APPLE, CRABAPPLE, PEAR, QUINCE	Do not apply within 30 days of harvest. **Dormant:** Overwintering Aphid and Mite Eggs, where shown: Putnam, San Jose, Scales — Use 2 pints with thorough coverage.
APRICOT: **Summer:** Codling Moth, Mites, Brown Mite, Two-Spotted — Use ½ pint. Do not apply within 30 days of harvest. 10 pints. **Dormant:** Overwintering Mite Eggs, Aphids, Peach, San Jose, Soft Brown — Use ½ to 1 pint.
CHERRY: **Summer:** Mites, Cherry Fruit, Clover, Two Spotted — Use ½ pint. Do not apply within 30 days of harvest. 10 pints. **Dormant:** Overwintering Aphid and Mite Eggs, Black Cherry, Forbes, Green, Parlatoria, San Jose, Scales — Use 2 pints.
CITRUS: Mites, [illegible]	[illegible] pints TRITHION 4-E per acre. Thorough application may be made up to 60 days before harvest. Do not apply within 14 days of harvest. Use not more than 10 pints of TRITHION 4-E per acre. If [illegible] not reapply within 30 days of the previous spray. Some State [illegible] pint TRITHION 4-E per acre. Do not apply within 30 days of harvest. 10 pints TRITHION 4-E per acre. Dormant and may be used in the regular summer oil [illegible] 41/2 pints TRITHION 4-E per acre. 10 pints TRITHION 4-E per acre. Do not apply within 30 days of harvest

...Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpen... ...Spotted Mite... Apply 1 pint. Do not apply within ... ...not make more than two applications per season. D... ...hay within 21 days after application.

...Aphid, Mexican Bean Beetle, Potato Leafhopper, Serpen... ...Spotted Mite... Apply 1 pint per acre. Do not make ... ...fications per season. Do not graze, cut for forage or ... ...n 21 days following the last application.

...an Aphid, Potato Flea Beetle, Serpentine Leaf Miner... ...Use 1½ to 1 pint. Do not apply within 21 days of harvest ...ed as feed or forage.

...RMELONS: Melon Aphid, Potato Flea Beetle, Serpentine ...otted Mite... Use 1 to 2 pints. Do not apply within 5 ...

...et): Grass Mite, Pacific Mite, Two Spotted Mite... Apply ...e more than one application per season. Do not graze or ... 21 days after application.

...n – Carmine, Cinnabar Mite, Two Spotted Mite, Tropical ... pint. **Mid and Late Season**... Use ¾ to 1 pint. If mite ..., use 1 to 1¼ pints and repeat in 5 to 10 days. Cotton ...o 1 pint. Cotton Aphid, Leaf Perforator, Leafhopper... Use ...ated plants should not be grazed or fed to livestock.

...ER SQUASH: Melon Aphid, Potato Leafhoppers, Serpen... ...-Spotted Mite... Use 1 to 2 pts. Do not apply within ...

...PIMENTO: Mites, Tarnished, Two Spotted, Potato Flea ...hopper, Serpentine Leaf Miner... Use 1 to 2 pints. D... ...ays of harvest.

...hopper, Two Spotted Mite... Use 1 pint. Do not apply ... ...harvest. Do not apply to ... ...olden Muskat, Muscat and the ...

...Sets or Transplants (for Dry Bulb or Green): ... ...Furrow treatment. **Muck Soils:** ... ...pints per acre. Apply at time of seeding or transplanting ...plant in furrow. Mix in 100 gallons of water and apply ...per 1,000 ft. of furrow. Keep the mixture agitated.

...rpentine Leaf Miner, Two-Spotted Mite... Use 1 to 1½ ... ...within 7 days of harvest. Do not feed treated forage ...

...Clover Mite, Cucumber Beetle, Sugarbeet Root Aphid, Sugarbeets **(listed on this label):** Aphids, Beet Armyworm, Beet Webworm, Spotted Wilt, Two Spotted Spider Mites, Western Yellow Striped Armyworm... Use 1 to 2 ... ...or feed crop residue to livestock. Sugarbeet seed crops ...o 14 days of harvest and crop residue can be fed t...

...g (Texas only), Pea Aphid, Sorghum Midge, Two Spotted ...Mite Species... 1 pint. Limit applications to two per ... or cut for forage within 21 days. Under certain environmental ...me varieties of grain sorghum, organophosphorus chemi... ...ght discoloration to the leaves.

...hid, Mexican Bean Beetle, Potato Leafhopper, Serpentine ...tted Mite. Apply 1 to 1¼ pints. Do not apply within 2 ...not feed treated forage to livestock.

...d, Serpentine Leaf Miner, Two-Spotted Mite... Use 1 ...ly within 21 days of harvest.

...berry Aphid, Two-Spotted Mite... Use 1 pint. Do not ... ...of harvest.

...Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two... 1 to 2 pints. Do not apply within 14 days of harvest.

...ato Russet, Tarnished, Two-Spotted, Potato Flea Beetle, **Potato Psyllid, Serpentine Leaf Miner**... Use 1 to ... ...within 7 days of harvest.

Stauffer

TRITHION® 4-E

EMULSIFIABLE LIQUID INSECTICIDE-ACARICIDE

CONTAINS FOUR POUNDS TRITHION PER GALLON

ACTIVE INGREDIENTS:	
Carbophenothion S-[(p-chlorophenylthio)methyl] O,O-diethyl phosphorodithioate	45.9%
Xylene Range Aromatic Hydrocarbon Solvent	45.5%
INERT INGREDIENTS:	8.6%
	100.0%

Rinse spray equipment and containers. Discharge liquid waste in pit in isolated area located away from water supplies.

PERFORATE EMPTY METAL CONTAINERS. CRUSH AND BURY GLASS.

U.S. Patent No. 2,793,224

WARNING

Keep Out of Reach of Children

MAY BE FATAL IF SWALLOWED.

Do not get in eyes, on skin or clothing. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. In case of contact, wash immediately with soap and water. For eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating, or smoking. Do not contaminate feed and food. Wash contaminated clothing with soap and hot water before re-use.

ANTIDOTE — Exposure may cause cholinesterase inhibition. Call a physician immediately; in all cases of suspected poisoning. Atropine is antidote. 2-PAM is also of benefit when administered early, and in conjunction with atropine.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Protect from temperatures below 0° F. Liquid phases may separate at lower temperatures. Warm or store at higher temperatures and mix thoroughly to recombine liquid phases and assure uniformity before use.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

COMBUSTIBLE! Do Not Use or Store Near Heat or Open Flame.

LOT NO. **GALLONS NET**

EPA REG. NO. 476-1376-AA

Made In U.S.A. By 2 ARC '26-23-

STAUFFER CHEMICAL COMPANY

WESTPORT, CT 06880

FRUIT AND NUT CROPS

...unless otherwise specified, dilute TRITHION 4 E and gallons spray for above shown **per 100 gallons water** for use ... thorough coverage spray. When used concentrate and semi-concentrate ... should be applied in sufficient water for thorough coverage according ... equipment manufacturers' directions.

...the maximum permissible pesticide rate per acre expressed as TRITHION ... E ... (Xylene or parenthesis ... after each ... and must not be ... regardless whether using dilute ... concentrate sprays. Apply a minimum of ... gallons TRITHION 4 E per acre ... from ... lower **minimum** ... after rate selected.

CROP	Gal. Water Per Acre Aircraft	Ground
Orchard, Mature Trees	10	40
Citrus	10	100

Almond (California only): Pre-Bloom or Petal-Fall: ... ...spray 1 pint. **Summer:** Mites... ... Use ½ pint plus ½ to 2 gallons spray oil. If almond hulls are to be used for feed, do not apply within 7 days of harvest or make more than one application per season. Multiple applications in accordance with local programs may be made if hulls are not to be used for feed. 5 pints. **Dormant:** Overwintering Mite and Aphid Eggs, Scales (European Fruit Lecanium, Olive (Parlatoria), San Jose... Use ½ to 1 pint plus 1 to 2 gallons spray ...

APPLE, CRABAPPLE, PEAR, QUINCE: Summer... ... Use ½ to 1. Apply only after full leaf expansion. Do not apply within 30 days of harvest. 2½ pints. **Dormant or Delayed Dormant:** Overwintering Aphid and Mite Eggs, Pear Psylla, Scales, Forbes, Oyster Shell, Putnam, San Jose, Scurfy... Use ½ to 1 pint plus 1 to 2 gallons spray oil with thorough coverage.

APRICOT: Summer: Codling Moth, Mites (Clover, Brown Almond), European Red, Two-Spotted... Use ½ pint. Do not apply within 30 days of harvest. 4 pints. **Dormant:** Overwintering Mite Eggs, Scales (European Fruit Lecanium, San Jose, Soft Brown... Use ½ to 1 pint plus 1 to 2 gallons spray oil.

CHERRY: Summer: Mites (Cherry Rust, Clover, Brown Almond), European Red, Two-Spotted... Use ½ pint. Do not apply within 30 days of harvest. 4 pints. **Dormant:** Overwintering Aphid and Mite Eggs, Scales, European Fruit Lecanium, Forbes, Olive (Parlatoria), San Jose... Use ½ to 1 pint plus 1 to 2 gallons spray oil.

CITRUS: ... ... Apply ½ to ¾ pints TRITHION 4 E per 100 gallons water. ½ pint is used, application may be made up to day of harvest. ¾ pints of ½ pint is used, do not apply within 11 days of harvest. ¾ pints. Do not apply more than 1,000 gallons of spray per acre. If more than one application is made, do not reapply within 30 days of the previous application. **Scales** (Florida only... Snow Scale... Use ½ pint TRITHION 4 E per 100 gallons water. Do not apply within 30 days of harvest. ¾10 pints TRITHION is compatible with oil and may be used in the regular summer oil spray program. EC 435-66 in summer, EC 412-66 in fall... 10 pints. TRITHION may injure grapefruit or Florida if combined with oil in the summer or if applied alone in the fall before the fruit is fully colored. Do not apply to Kumquat or Citron. Scales, California Red, Cottony, Cushion, Florida Red, Purple, Brown, Yellow... ½ pint plus 1 to 2 gallons spray oil. Do not apply within 30 days of harvest. ¾10 pints. TRITHION may cause off-colored spots on grapefruit in Texas if applied after July 1 and before fruit is fully colored or in summer or fall before fruit is fully colored in other areas. Do not apply to Kumquat or Citron.

FIGS: ... ... Mite... Use ½ to 1 pint. Do not apply with a 7 day ... harvest. ... pints.

OLIVE: Scales, Black, Citricola, ... Parlatoria, ... San Jose... Apply ... pint plus ... gallons of spray oil. Do not use within ... of ... ½ pint.

PEACH, NECTARINE: Summer: Mites, Clover, Brown ... Two Spotted... Use ½ pint. Do not apply within ... **Dormant:** Scales (European Fruit Lecanium, Olive, Parlatoria, San Jose... Use ½ to 1 pint plus 1 to 2 gallons spray oil. Overwintering Mite Eggs, normal only... Use 1 pint.

PECAN: Two-Spotted Mite... Use ½ pint. Do not apply after husks open. Do not graze livestock... Cover crops grown in treated orchards may be grazed.

PLUM, PRUNE: Summer: Leaf Curl Plum Aphid, Leafhoppers, Mites (Brown Almond, European Red, Two Spotted... Use ½ pint. Do not apply within 30 days of harvest. 4 pints. **Dormant:** Overwintering Aphid and Mite Eggs, Scales (European Fruit Lecanium, Olive, Parlatoria, San Jose... Use ½ to 1 pint plus 1 to 2 gallons spray oil.

WALNUT: Summer: Codling Moth, European Red Mite, Two Spotted Mite, Walnut Aphid, and for suppression of European Fruit Lecanium... Use ½ pints per acre in sufficient water for coverage. Best results for scale insect suppression are obtained if application is made after eggs have hatched. Do not apply after husks open. Do not graze livestock in cover crops in treated Walnut groves. **Dormant:** Walnut Aphid, Overwintering Mite Eggs, Scales (Calico, European Fruit Lecanium, Frosted... Use ½ to ¾ pints plus 1 gallons spray oil per acre in sufficient water for coverage. Do not use more than 2 gallons of oil per acre.

POST HARVEST APPLICATION TO DECIDUOUS FRUIT AND NUT CROPS: To ... ... overwintering populations of Aphids, Leafhoppers, Spider Mites... ... ½ to 1 pint Trithion 4-E ... control of Aphid and Mite Eggs, Scales, European Fruit Lecanium, San Jose... ... ½ to 1 pint Trithion 4-E plus 1 to 2 gallons spray oil. Cover crops should not be grazed or fed to livestock.

NON-FOOD CROPS (For use by commercial applicators only)

FOREST AND CHRISTMAS TREE PLANTINGS: ... ... Apply 1½ to 2 pints ... 5 to 5... gallons water for acre. Repeat application ... necessary.

...TURF: Chinch Bug, Bermuda Mite, Rhodes Grass Scale, and Thatch in Bermudagrass – Wet ... thoroughly, then apply 12½ pints of TRITHION 4 E in 100 to 150 gallons water to 5,000 sq. ft. of lawn area. Do not permit children or domestic animals on lawn until treated area is thoroughly dry.

ORNAMENTALS: Arbor-Vitae, Azalea, Hollyhock, Aphids... Asparagus Fern, Begonia, Boxwood, Cactus, Caladium, Chrysanthemum, Cineraria, Citrus, Croton, Coleus, Cosmos, Cotoneaster, Crataegus, Hawthorn, Cycas, Sago-Palm, Dahlia, Delphinium, Eucalyptus, Euonymus, Euphorbia pulcherrima (Poinsettia), Fagus, Beech, Ficus Elastica (Rubber Plant), Ferns, Firs, Fuchsia, Gardenia, Guava, Hedera Helix (Ivy), Hemlock, Hydrangea, Ilex (Holly), Jasmine, Juniper, Lantana, Lathyrus (Sweet Pea), Ligustrum (Privet), Lilium (Lily), Magnolia, Myrtus (Myrtle), Nerium (Oleander), Orchids, Palms, Phlox, Pine, Pittosporum, Sycamore, Poplar, Primula (Primrose), Prunus, Almonds, Peaches, Cherries, Plums, Pyracantha (Firethorn), Rhododendron and Azalea, Ribes (Flowering Currant), Robinia, Black Locust, Saintpaulia (African Violet), Sorbus (Mountain Ash), Strelitzia (Bird-of-Paradise), Syringa (Lilac), Tilia (Linden), Tropaeolum (Nasturtium), Ulmus (Elm), Violas (Violets), Pansies.

For Control of: Aphids (Monterey Pine, Rose, Black Bean, Pine Bark, Apple, Rosy Apple, Woolly Apple, Oleander, Mexican), ... (Citrus Red, Florida Red, Purple, Brown, Yellow), ... Yuma, Scale Insects, Black, Hemispherical, Latania, Soft, California Red, ... Cotton, Cushion, Yellow, ... Pine, ... Pear, ... Peaches, Bayberry, ... to 2 pints... ... gallons of water.

CAUTION: Tolerance of most plants to TRITHION appears to be excellent. A few ornamentals and old or some particular certain forms and varieties are sensitive; greenhouse roses have been injured. When ... of ... TRITHION, ... ½ pints. Use ... or ... pints.