

50 LB.

Apr 1964



NET

SEVIN* 5 DUST

AN AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENT:	Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS		95.0%
		100.0%

*Trade Mark of Stauffer Chemical Company for the active ingredient, 1-naphthyl N-methylcarbamate, Carbaryl.

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED OR INHALED.

- Avoid prolonged or repeated breathing of dust.
- Avoid prolonged or repeated contact with skin.
- Skin Contact with Any Insecticide May Be Harmful.
- Wash hands and face before eating or smoking. Take shower or bath after work.
- Wear regular long-sleeved clothing. Change to clean clothing daily.
- Avoid contamination of food, feed, feeding troughs and watering receptacles.
- Note for Physicians: Carbaryl is a carbamate. Avoid high blood pressure, asthma, and other conditions.
- Avoid eye contact.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Apply this insecticide to crops and plants only. Do not use on animals or humans. Do not use on lawns, lawns, or ornamentals. Do not use on watermelons in Florida. Do not use on strawberries within 1 day of harvest. Do not use on tobacco within 1 day of harvest. Do not use on corn within 14 days of harvest. Do not use on carrots, Chinese cabbage, collards, endive, escarole, kale, lettuce, mustard greens, parsley, parsnips, radishes, rutabagas, spinach, Swiss chard, turnips, and other leafy vegetables within 3 days of harvest. Do not use on beans, broccolis, Brussels sprouts, cabbages, cauliflowers, eggplants, peppers, potatoes, tomatoes, and other vegetables within 14 days of harvest. Do not use on corn, sorghum, and other grain crops within 14 days of harvest. Do not use on alfalfa, clovers, cowpeas, and other legume crops. Do not use on cotton, flax, hemp, and other fiber crops. Do not use on citrus fruits, grapes, and other fruit crops. Do not use on ornamentals, lawns, and lawns. Do not use on watermelons in Florida. Do not use on strawberries within 1 day of harvest. Do not use on tobacco within 1 day of harvest. Do not use on corn within 14 days of harvest. Do not use on carrots, Chinese cabbage, collards, endive, escarole, kale, lettuce, mustard greens, parsley, parsnips, radishes, rutabagas, spinach, Swiss chard, turnips, and other leafy vegetables within 3 days of harvest. Do not use on beans, broccolis, Brussels sprouts, cabbages, cauliflowers, eggplants, peppers, potatoes, tomatoes, and other vegetables within 14 days of harvest. Do not use on corn, sorghum, and other grain crops within 14 days of harvest. Do not use on alfalfa, clovers, cowpeas, and other legume crops. Do not use on cotton, flax, hemp, and other fiber crops. Do not use on citrus fruits, grapes, and other fruit crops.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when insects are actively present or about to appear. Unless otherwise specified, apply at the first sign of infestation and repeat at 7-10 day intervals when needed to maintain control, but do not exceed the maximum number of applications specified.

RECOMMENDATIONS

- ALFALFA, CLOVERS, COWPEAS** Southern Peas, PEANUTS, SOYBEANS Armyworms, Corn Earworms, Stinkbugs, Webworms - Use 2-30 lb. per acre. Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetles, Green Cloverworm, Japanese Beetle, Leafhopper, Three-cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar - Use 40 lb. per acre. Grasshoppers - Use 10-30 lb. per acre. Cowpea Curculio - Use 40 lb. per acre.
- BEANS** Mexican Bean Beetle - Use 17 lb. per acre. Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvetbean Caterpillar, Western Bean Cutworm - Use 40 lb. per acre. Armyworms, Corn Earworm, Stinkbugs, Tarnished Plant Bugs - Use 20-30 lb. per acre.
- BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER** Flea Beetle, Harlequin Bug - Use 2-30 lb. per acre. Armyworms, Corn Earworm, Cabbage Caterpillars - Use 40 lb. per acre. Do not apply within 3 days of harvest.
- CARROTS, CHINESE CABBAGE, COLLARDS, ENDIVE, ESCAROLE, KALE, LETTUCE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SPINACH, SWISS CHARD, TURNIPS** Flea Beetles, Harlequin Bug, Leafhoppers - Use 10-20 lb. per acre. Six-spotted Leafhopper - Use 40 lb. per acre. Armyworms, Cabbage Caterpillars, Corn Earworm, Stinkbugs, Tarnished Plant Bugs - Use 20-30 lb. per acre. Do not apply within 3 days of harvest on head lettuce, parsnips, radishes, rutabagas, and turnips; 14 days of harvest if tops are to be used as food or feed. Do not apply within 14 days of harvest on leaf lettuce and other leaf crops.
- CORN** Corn Earworm, Corn Rootworm, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhoppers, Sap Beetles - Use 20-40 lb. per acre. Do not use on corn within 14 days of harvest.
- CUCUMBER, MELONS, PUMPKIN, SQUASH** Cucumber Beetle, Flea Beetles, Leafhoppers, Squash Bug - Use 10-20 lb. per acre. Melonworm, Pickleworm - Use 40 lb. per acre. Do not use on watermelons in Florida. May be applied up to 14 days including the day of harvest.
- EGGPLANT, PEPPER, POTATO, TOMATO** Colorado Potato Beetle, Flea Beetles, Leafhoppers - Use 10-20 lb. per acre. European Corn Borer, Fall Armyworm, Flea Beetles, Stinkbugs, Tarnished Plant Bugs, Tomato Fruitworm, Tomato Hornworms - Use 20-40 lb. per acre. May be applied up to day of harvest.
- OKRA** Corn Earworm, Stinkbugs - Use 20-30 lb. per acre.
- TOBACCO** In Plant Bug, Tobacco Fly, Beetle - Use 20-40 lb. per acre. Do not use on tobacco within 1 day of harvest.
- STRAWBERRIES** Leaf Roller, Meadow Spittlebug, Strawberry Weevil - Use 40 lb. per acre. Do not apply within 1 day of harvest.

BURN EMPTY CONTAINER - KEEP OUT OF SMOKE.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.