


# CAPTAN-PCNB 10-10 DUST



## SOIL FUNGICIDE SEED PROTECTANT

ACTIVE INGREDIENTS	
CAPTAN: N[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide .....	10%
Pentachloronitrobenzene .....	10%
INERT INGREDIENTS: .....	80%
	100%

U. S. Pat. Nos. 2,553,770—2,553,771  
Not licensed for use or export outside of the United States, its territories and possessions, Canada or Mexico

### CAUTION

Keep Out of Reach of Children

#### HARMFUL IF SWALLOWED

MAY CAUSE SKIN IRRITATION.  
AVOID BREATHING DUST, CONTACT WITH EYES, SKIN AND CLOTHING  
WEAR RESPIRATOR AND GOGGLES.  
WASH THOROUGHLY AFTER HANDLING.  
AVOID CONTAMINATION OF FEED & FOODSTUFFS.

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT.

#### USE PRECAUTIONS

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

**This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of dust from areas treated.**

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-1500  
S-SF-680806-WC

Made in U.S.A.  
Stauffer Chemical Company  
New York, N.Y.

50 LB. NET

#### RECOMMENDATIONS

CAPTAN-PCNB 10-10 D U S T is a special formulation for use at time of planting to protect beans and cotton seed and seedlings against soil borne diseases.

#### DIRECTIONS FOR USE

##### COTTON

For protection against damp-off, seed rot,  
Rhizoctonia (Soreskin) and Phythium.

**IN-FURROW APPLICATION:** Apply 15 pounds of Captan-PCNB 10-10 D u s t per acre based on a 30" row spacing. Increase the rate per acre proportionately as the row spacing decreases. For example, on a 24" row spacing, use 18¾ pounds per acre; on an 18" row spacing, use 25 pounds per acre, and on a 14" row spacing, use 32 pounds per acre. Application should be made to the planting furrow and covering soil at the time of planting.

**PLANTER BOX APPLICATION:** Use 5 pounds Captan-PCNB 10-10 D u s t per acre. The seed and the proper amount of this material may be pre-mixed in a container just prior to dumping in the planter box, or the chemical may be added to the seed in the planter box and blended by stirring or mixing thoroughly. Planter box application is not intended as a substitute for regular seed treatment. Is a supplement to proper chemical treatment of good seed.

**HILL-DROP SEEDING — PLANTER BOX APPLICATION:** Use 3 pounds Captan-PCNB 10-10 D u s t for hill-drop seeding rate of 15-25 pounds per acre. Balance of directions for planter box application given above apply to hill-drop seeding method.

##### BEANS

**IN-FURROW APPLICATION:** For protection against root rot (Rhizoctonia) and stem rot (Pythium), apply 10-15 pounds Captan-PCNB 10-10 D u s t per acre based on a 30" row spacing. Increase rates per acre proportionately as the row spacing decreases. Application should be made to the planting furrow and covering soil at time of planting.

**NOTE:** To insure correct planting rate, calibrate planter for proper seeding rate after adding the recommended dosage of Captan-PCNB 10-10 D u s t.

**BURN EMPTY BAG—KEEP OUT OF SMOKE**