

476-1514
ACCEPTED
 Jan 21, 1971
 UNDER THE FEDERAL INSECTICIDE
 ACT AND REGULATIONS THEREUNDER
 THE FOLLOWING PESTICIDE
 IS REGISTERED FOR SALE AND
 DISTRIBUTION IN THE UNITED STATES
 OF AMERICA



CAPTAN-HEXACHLORO BENZENE 40-40 SEED PROTECTANT

USDA Reg. No. 476-1514

ACTIVE INGREDIENTS

CAPTAN: N-[[trichloromethyl]thio]-4-cyclohexene-1,2-dicarboximide 40%
 Hexachlorobenzene 40%

INERT INGREDIENTS: 20%
 100%

*Not licensed for use in countries where valid patents exist

CAUTION Keep Out of Reach of Children
 HARMFUL IF TAKEN INTERNALLY

AVOID INHALING DUST. WEAR RESPIRATOR.
 AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
 WASH WITH SOAP AND WATER AFTER HANDLING.
 AVOID CONTAMINATION OF FEED AND FOOD.

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT.

DESTROY EMPTY CONTAINERS - BURY OR BURN -
 STAY OUT OF SMOKE AND FUMES

NET WT. LBS.

STAUFFER CHEMICAL COMPANY
 NEW YORK, N.Y. 10017

USE PRECAUTIONS

Toxic to fish. Do not apply where any body of water may be contaminated by direct application or runoff. Treated seeds exposed on the soil surface will be hazardous to birds and other wildlife.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

IMPORTANT: Treated seed must not be used for human consumption, nor as animal feed. Treated seed should be labeled:

"TREATED SEED — DO NOT USE FOR FEED, FOOD OR OIL PURPOSES."

The Federal Government and some States have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed. Information on this subject may be obtained from local authorities.

CAPTAN-HEXACHLOROBENZENE 40-40 SEED PROTECTANT is a special formulation for the treatment of WHEAT SEEDS prior to planting for the protection of the seeds against Bunt (Stinking Smut), seed rots, seedling blights and root rot.

CAPTAN-HEXACHLOROBENZENE 40-40 SEED PROTECTANT IS A VERSATILE FORMULATION WHICH CAN BE USED TO TREAT WHEAT SEED BY THE SLURRY METHOD, PLANTER BOX METHOD OR DUST TREATMENT.

DIRECTIONS FOR USE

PLANTER BOX TREATMENT - 1 oz. per bushel seed (1 2/3 ozs. per 100 lbs. seed). Mix thoroughly with seed in planter box before planting.

DUST TREATMENT - Apply 1 ounce per bushel of seeds (1 2/3 ozs. per 100 lbs. seed). Mix thoroughly with seed in dusting machine, barrel or other container. May also be applied through drill box or planter box.

SLURRY TREATMENT - Apply at the rate of 1 ounce per bushel. For 46 cc cup, using dump weight of 10 lbs., add 1 lb. CAPTAN-HEXACHLOROBENZENE 40-40 SEED PROTECTANT to 1 gallon water. For 23 cc cup, using dump weight of 10 lbs., add 2 lbs. CAPTAN-HEXACHLOROBENZENE 40-40 SEED PROTECTANT to 1 gallon water. For "mist type" cup setting of 15, set machine to dump 6.6 lbs., add 2 lbs. CAPTAN-HEXACHLOROBENZENE 40-40 SEED PROTECTANT to 1 gallon water. One gallon of slurry will treat approximately 1650 lbs. of seed.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

S-SF-7000 (1-11)