



INDICATIONS  
FRUIT TREES  
Sprays



GAL. NET

# TRITHION<sup>®</sup> 6-E

**EMULSIFIABLE LIQUID  
AN ORGANIC PHOSPHATE INSECTICIDE-  
ACARICIDE**

**ACTIVE INGREDIENT:**

CARBOPHENOTHION S-[[[p-chlorophenyl]thio] methyl]  
O,O-diethyl phosphorodithioate

Xylene range aromatic solvent

**INERT INGREDIENTS:**

62.3%  
22.0%  
15.7%  
100.0%

TRADE MARK OF STAUFFER CHEMICAL CO.  
U.S. PAT. NO. 2,793,224

## WARNING

**Keep Out of Reach of Children**

**MAY BE FATAL IF SWALLOWED**

**DO NOT GET IN EYES, ON SKIN OR CLOTHING.** If in eyes, flush with water for 15 minutes. If on skin, wash immediately with soap and water. If on clothing, wash with water. If on hands, arms and face, wash with soap and water for 15 minutes. If on clothing, wash with water.

**ATROPINE IS ANTIDOTAL.** Symptoms of atropine poisoning include dry mouth, blurred vision, and difficulty urinating. If these symptoms occur, give atropine immediately. For information on atropine, contact your doctor or pharmacist.

ESTROY EMPTY CONTAINERS IMMEDIATELY  
**STAUFFER CHEMICAL CO.**  
NEW YORK

## RECOMMENDATIONS Cont

### FIELD CROPS

Use rate late evening or early morning. Apply with sufficient water to provide adequate coverage.

**ALFALFA:** Pea aphid, potato leafhopper, alfalfa weevil, two spotted mite. — Apply 11.5 pts. per acre. Do not apply more than 1 application per season. Do not graze or cut hay within 21 days after application.

**BEANS, DRY (Shelled):** Bean aphid, alfalfa weevil, leafhopper, serpentine leaf miner, two spotted mite. — Apply 11.5 pts. per acre. Do not apply more than 1 application per season. Do not graze or cut hay within 21 days after application.

**BEANS, SNAP:** Bean aphid, alfalfa weevil, leafhopper, serpentine leaf miner, two spotted mite. — Apply 11.5 pts. per acre. Do not apply more than 1 application per season. Do not graze or cut hay within 21 days after application.

**CLOVER:** Pea aphid, potato leafhopper, alfalfa weevil. — Apply 11.5 pts. per acre. Do not apply more than 1 application per season. Do not graze or cut hay within 21 days after application.

**CORN (Field & Sweet):** Corn earworm, European corn borer, fall armyworm, European spruce sawfly. — Apply 11.5 pts. per acre. Do not graze or cut hay within 21 days after application.

**COTTON, SOUTH ONLY (Early Season):** Cotton leaf miner, two spotted mite, tobacco budworm. — Apply 11.5 pts. per acre. Do not graze or cut hay within 21 days after application.

**(Mid and Late Season):** Cotton leaf miner, two spotted mite, tobacco budworm. — Apply 11.5 pts. per acre. Do not graze or cut hay within 21 days after application.

**OTHER AREAS:** For control of aphid, spider mites, and other insects on ornamentals, apply 11.5 pts. per acre. Do not apply more than 1 application per season.

**SORGHUM:** Two spotted mite, European spruce sawfly. — Apply 11.5 pts. per acre. Do not graze or cut hay within 21 days after application.

### NON-FOOD CROPS

For Use by Commercial Applicators Only

**PEANUTS:** Peanut aphid, peanut root-knot disease. — Apply 11.5 pts. per acre. Do not apply more than 1 application per season.

**ORNAMENTALS:** For control of aphid, spider mites, and other insects on ornamentals, apply 11.5 pts. per acre. Do not apply more than 1 application per season.

**For Control of Aphid:** Apply 11.5 pts. per acre. Do not apply more than 1 application per season.

**For Control of Spider Mites:** Apply 11.5 pts. per acre. Do not apply more than 1 application per season.

**CAUTION:** Do not apply to plants that are stressed or damaged. Do not apply to plants that are in bloom.

1951-916-1591  
6-16-67