

10-13-71
415-1612

DIRECTION FOR USE

GAL. NET

LINDANE

AN AGRICULTURAL INSECTICIDE
AN EMULSIFIABLE LIQUID

1-EH

Contains 1 lb. Lindane Per Gallon

ACTIVE INGREDIENTS:

Lindane 1.00%
Emulsifier 0.50%
Inert Ingredients 98.50%

INERT INGREDIENTS:

CAUTION

Keep Out of Reach of Children

POISONOUS TO MAN AND ANIMALS.
Irritates eyes, nose, throat, and skin.
Avoid breathing vapors.
Avoid contact with skin.
Do not use or store near heat or open flame.
Do not use near food or animal feed.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

OBSERVE THE FOLLOWING USE PRECAUTIONS:
FOR USE ONLY ON CROPS RECOMMENDED ON THIS LABEL.

Always
Effort

RECOMMENDED
RATES GIVEN BELOW
OF THIS MATERIAL PER
FOR THOROUGH

BRASSICAE - Broccoli, Brussels
Sprouts, Cabbage, Cauliflower, Kale,
Kohlrabi, Turnip, Turnip Greens

CUCURBITS - Cucumber, Melon, Pumpkin,
Squash, Zucchini

TOMATOES - Greenhouse and Field

FLY C

GENERAL INFORMATION - RESIDUE
Lindane is a contact insecticide.
It is effective against many insects.
It is not effective against sucking insects.
It is not effective against mites.

DO NOT USE NEAR MANUFACTURED
APPLIANCE - Do not use
near electrical appliances or
other equipment.

10-13-71
10-16-72

DIRECTIONS FOR USE

1 GAL NET

LINDANE

AN AGRICULTURAL INSECTICIDE

1-EH

AN EMULSIFIABLE LIQUID

Contains 1 lb. Lindane Per Gallon

ACTIVE INGREDIENTS:

gamma-hexachlorocyclopentadiene	1.0
Emulsifier	3.1
Emulsifier	71.2
INERT INGREDIENTS:	3.6
<hr/>	
100.0	

CAUTION

Keep Out of Reach of Children

HARMFUL IF SWALLOWED
 MAY BE ABSORBED THROUGH SKIN
 AVOID BREATHING SPRAY MIST
 AVOID CONTACT WITH SKIN
 IN CASE OF CONTACT, WASH IMMEDIATELY WITH LARGE AMOUNTS OF WATER
 KEEP OUT OF REACH OF CHILDREN AND PETS
 KEEP OUT OF REACH OF CHILDREN AND PETS
 DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

RECOMMENDATIONS
 DOSAGE RATES GIVEN BELOW ARE IN TERMS OF QUART
 OF THIS MATERIAL PER ACRE IN SUFFICIENT WATER
 FOR THOROUGH COVERAGE.

CUCURBITS (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Aphids, Trips) Use 1/2 quart per acre. Do not apply after edible parts start to form.

CUCURBITS (Cantaloupe, Cucumber, Melons, Squash) Aphids, Leaf Miners, Mealy Worm, Trips Use 1/2 quart per acre. Do not apply within 7 days of harvest in cucumbers and squash.

TOMATOES: Aphids, Colorado Potato Beetles, Leaf Miners, Trips Use 1/2 quart per acre. Do not use after fruit starts to form.

FLY CONTROL

GENERAL FLY CONTROL RESIDUAL SPRAY: Use 1/2 quart per acre. Do not use 2-4 gallons per acre. Spray as a dust or spray thoroughly wet walls, windows, doors, and other places where flies rest or congregate. Never apply so heavily that spray runs in places. Repeat application when fly control diminishes.

NOTICE: The manufacturer makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose concerning this material, except those which are contained on the main factory label.