

OBSERVE THE FOLLOWING USE PRECAUTIONS

Stauffer
CHEMICALS

1 GAL. NET

VAPONA*

INSECTICIDE

1.4 E

DAIRY & LIVESTOCK SPRAY
DDVP EMULSIFIABLE CONCENTRATE

For Controlling Face Flies and Certain Other Livestock Pests

Contains 1.4 lbs. VAPONA Insecticide per Gallon



ACTIVE INGREDIENTS	
2,2-Dichloro-1,1-Dimethyl-Phosphate	17.2%
Resistant Compounds	1.3%
Xylene Range Aromatic Hydrocarbon Solvent	79.5%
INERT INGREDIENTS	
	11.0%
	100.0%

Patented in U.S. VAPONA Insecticide
SHELL REGISTERED TRADEMARK



DANGER -- POISON



Keep Out of Reach of Children

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES
Wash thoroughly with soap and water. Do not eat, drink, or use tobacco products while using this product. Avoid contact with eyes, nose, mouth, and skin. If contact occurs, wash immediately with plenty of water. Do not breathe vapors. Do not use in confined spaces. Do not use near open flame or heat. Do not use in areas where food is prepared or stored. Do not use in areas where children play. Do not use in areas where livestock are kept. Do not use in areas where crops are raised. Do not use in areas where water is used for drinking. Do not use in areas where water is used for irrigation. Do not use in areas where water is used for other purposes. Do not use in areas where water is used for other purposes.

BEFORE USING, STUDY POISONING SYMPTOMS, FIRST AID TREATMENT AND
ANTIDOTE GIVEN ON LEFT PANEL OF THIS LABEL

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME
PROTECT FROM TEMPERATURES BELOW 0°F

nor crop lands

STAUFFER CHEMICAL CO.

NEWARK

POISON

ANTIDOTE AND FIRST AID TREATMENT

IF SWALLOWED: Do not induce vomiting. Give 1-2 glasses of water. If unconscious, give water by mouth. If vomiting occurs, do not give more water. If breathing is difficult, give oxygen. If necessary, give artificial respiration. If unconscious, give water by mouth. If vomiting occurs, do not give more water. If breathing is difficult, give oxygen. If necessary, give artificial respiration.

SEE LEFT PANEL



1 GAL. NET

VAPONA*

INSECTICIDE

1.4 E

DAIRY & LIVESTOCK SPRAY DDVP EMULSIFIABLE CONCENTRATE

For Controlling Face Flies and Certain Other Livestock Pests

Contains 1.4 lbs. VAPONA Insecticide per Gallon



ACTIVE INGREDIENTS:	
2,2-Dichlorovinyl Dimethyl Phosphate	17.2%
Related Compounds	1.3%
Xylene Range Aromatic Hydrocarbon Solvent	70.5%
INERT INGREDIENTS	
	11.0%
	100.0%

Patent No. 1851 VAPONA Insecticide
SHELL REGISTERED TRADEMARK



DANGER - POISON



Keep Out of Reach of Children

POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN OR EYES.

Wear protective suit or clothing. Do not breathe fumes. Avoid exposure to fumes. Wear protective clothing and goggles when mixing the concentrate to prepare emulsion. Wear protective clothing and goggles when spraying. Registered under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, by the U.S. Department of Agriculture for DDVP, VAPONA 1.4 E. Keep out of reach of children and animals. In case of contact, wash with soap and water. Wash hands, arms, and face thoroughly with soap and water after use. Do not eat, drink, or smoke until thoroughly washed with water. If inhaled, get fresh air and water. If in eyes, flush with water.

BEFORE USING STUDY POISONING SYMPTOMS, FIRST AID TREATMENT AND ANTIDOTE GIVEN ON LEFT PANEL OF THIS LABEL.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

PROTECT FROM TEMPERATURES BELOW 0°F

Use on non-crop land only. Do not use on crops.

STAUFFER CHEMICAL CO.

NEW YORK

PRODUCT INFORMATION

VAPONA Insecticide 1.4 E Dairy & Livestock Spray Concentrate is a specially prepared animal insecticide for use in the control of certain insects attacking livestock (Face flies, Houseflies, Horn flies, Stable flies, mosquitoes and gnats).

MIXING DIRECTIONS

1% Solution: Mix 8 fl. ozs. (1 cup) concentrate with 1 gallon water.
0.5% Solution: Mix 4 fl. ozs. (1/2 cup) concentrate with 1 gallon water.

RECOMMENDATIONS

BEEF AND DAIRY CATTLE: Flies (Houseflies, Horn flies, Stable flies, Face flies) and Mosquitoes. Apply 1 to 2 ozs. of 1% solution per animal as a mist spray daily with a hand or automatic sprayer. Spray thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply in excess of 2 fluid ounces per animal per day.

Do not apply to animals which are pregnant or nursing. Do not apply to animals which are to be treated with any type of salt or mineral supplements.

Face Fly: Mix 1/2 cup of face fly concentrate with 1 to 2 cups of sugar to each gallon of 1% solution.

CATTLE FEED LOTS, STOCK YARDS, HOLDING PENS & CORRALS: Flies, Mosquitoes & Gnats. Apply at the rate of 4 to 3 gallons of 0.5% solution per acre with a mist sprayer or other suitable equipment as an overall application. The mist spray should be directed over the entire area. Particular attention should be given to areas where flies congregate such as fences, spillage areas around feed bunk and walls of buildings. Animals may be present during treatment. Avoid direct application to exposed feed and water. For baited solutions, dissolve one pound of sugar per 5 to 10 gallons of water before adding the Vapona concentrate.

ANIMAL BUILDINGS: Stables, Barns, Milk Sheds, Shelter Sheds, Horse Barns and Outdoor Areas. For control of Flies, gnats and mosquitoes. Before applying reduce air movement as much as possible by closing doors, windows and other openings.

Surface Spray: Apply 1/2 cup of concentrate to 1 gallon of water as a coarse wet spray per 1000 sq. feet of milk sheds, stables, livestock barns, milk sheds, poultry houses and other areas that are not dryways. Feed storage bins, a house and water troughs. **Poultry:** House apply to manure with a sprayer. Consider the legs, feet, sides and bottoms of feet, troughs and water troughs. Avoid contact with wings and beak of poultry. Repeat treatments as necessary to control flies or barns.

Fog or Mist: Use 1 cup of concentrate to 1 gallon of water to cut feet of flies. Animals should be kept in a dark place until they have been treated and treated again with high pressure.

CAUTION: Do not use in water troughs, feed troughs, or water troughs.

FLY MAGGOTS: Do not use on manure. Do not use on manure.

NOTICE: The manufacturer makes no warranty, express or implied, including the warranties of merchantability and or fitness for any particular purpose concerning the material, except those which are contained on the manufacturer's label.

476-1120