



ETHION 25-WP

ORGANOPHOSPHORUS INSECTICIDE-ACARICIDE WETTABLE POWDER - FOR USE IN WATER

ACTIVE INGREDIENT:

*Ethion (O,O,O',O'-tetraethyl S,S'-methylene bisphosphorodithioate)

INERT INGREDIENTS:

U.S. Patent No. 2,873,228

WARNING

**MAY BE FATAL IF SWALLOWED.
ABSORBED THROUGH SKIN.**

**DO NOT BREATHE VAPOR, DUST OR SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
WASH THOROUGHLY AFTER HANDLING.
DO NOT CONTAMINATE FOOD OR FEED.**

Keep Out of
Reach of Children

476-1792
ACCEPTED
Jan 16 1971

USE PRECAUTIONS

Observe use limitations on this material. Do not use on fish and wildlife. Keep out of any body of water. Apply this product only as directed on this label.

In order that pesticide residues on food and feed crops will not exceed tolerance levels, use the appropriate label and Do Not Exceed Rate of Application rates and intervals, and do not apply more than the specified harvest interval. Do not apply to areas where animals are present. The grower is responsible for any damage caused by drift from the property to that of others.

Consult state Agricultural Experiment Station or State Agricultural Extension Service for information on the toxicity of this material to animals. Do not use on animals.

This product is not to be used on any crop or animal which is listed on the label as being sensitive to this material.

BURN EMPTY BAG - KEEP OUT OF SMOKE

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

LB. NET

STAUFFER CHEMICAL COMPANY NEW YORK, N.Y.

ETHION 25-WP

DIRECTIONS FOR USE

Apply this material to the crop or animal as directed on the label. Do not use on fish and wildlife. Keep out of any body of water. Apply this product only as directed on this label.

RECOMMENDATIONS

ALMONDS: (Sweet and Dry) Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

25.0%
75.0%
100.0%

APPLES: (Sweet and Dry) Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

BEANS: (Succulent and Dry) Use 1 lb per acre. Do not apply more than 2 days before harvest. Do not apply within 30 days of harvest.

BEAN SEED TREATMENT: Use 1 lb per acre. Do not apply more than 2 days before harvest. Do not apply within 30 days of harvest.

CHERRIES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

CITRUS: (Citrus-fruit, orange, lemons, limes, tangelos, tangerines) Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

COTTON: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

EGGPLANTS: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

GRAPES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

MELONS: (Watermelon, cantaloupe, honeydew, muskmelon) Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

PEACHES, NECTARINES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

PEPPERS, PIMIENTOS: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

PLUMS, PRUNES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

STRAWBERRY: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

TOMATOES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

WALNUT: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

Directions for Use

Red Scale, Citrus Whitefly: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

California and Arizona: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

Do not apply more than 3 times per year. Do not apply within 21 days of harvest. Do not apply on tangerines, do not apply on crop. Do not use oil.

MELON: (Watermelon, cantaloupe, honeydew, muskmelon) Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

ONIONS: (Set, Onion) Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

PEACHES, NECTARINES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

NECTARINES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

PEPPERS, PIMIENTOS: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

PLUMS, PRUNES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

STRAWBERRY: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

TOMATOES: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.

WALNUT: Use 1 lb per acre. Do not apply more than 3 days before harvest. Do not apply within 30 days of harvest.



ETHION 25-WP

PHOSPHORUS INSECTICIDE-ACARICIDE WATER SOLUBLE POWDER—FOR USE IN WATER

25.0%	25.0%
75.0%	75.0%
100.0%	100.0%

U.S. Patent No. 2,873,278

WARNING

FATAL IF SWALLOWED.
POISONOUS IF INhaled THROUGH SKIN.

AVOID THE VAPOR, DUST OR SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
WASH THOROUGHLY AFTER HANDLING.
DO NOT CONTAMINATE FOOD OR FEED.

PRECAUTIONS

Always use in accordance with directions given on the label for the crop and area. Do not apply to fish and wildlife. Keep out of any body of water. Apply this product to crops and areas only when it is safe to do so. Do not exceed the rates established by the manufacturer. Do not apply at intervals shorter than those recommended. Do not apply to lawns, golf courses, pastures, or other areas where the general public has access. The grower is responsible for any damage caused by drift from his property to that of others.

State Agricultural Experiment Stations or State Agricultural Extension Services should be consulted for information on the use of this material. Do not use this material in a manner inconsistent with the label directions.

BURN EMPTY BAG — KEEP OUT OF SMOKE

Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and fitness for a particular purpose, concerning this material, except those which are contained on Stauffer's label.

1 LB. NET

STAUFFER CHEMICAL COMPANY NEW YORK, NY 10017

ETHION 25-WP DIRECTIONS FOR USE

RECOMMENDATIONS

ALMONDS: (Succulent and Dry) To control almond scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest.

APPLES: Not more than 1 application during fruiting period. Do not apply within 21 days of harvest. Do not apply to apples that are to be eaten raw. Do not apply to apples that are to be processed. Do not apply to apples that are to be stored.

BEANS: (Succulent and Dry) To control bean leaf beetle, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to beans that are to be eaten raw. Do not apply to beans that are to be processed. Do not apply to beans that are to be stored.

BEAN SEED TREATMENT: To control seed beetle, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to beans that are to be eaten raw. Do not apply to beans that are to be processed. Do not apply to beans that are to be stored.

CHERRIES: To control cherry scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to cherries that are to be eaten raw. Do not apply to cherries that are to be processed. Do not apply to cherries that are to be stored.

COTTON: To control cotton scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to cotton that is to be ginned. Do not apply to cotton that is to be stored.

EGGPLANTS: To control eggplant scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to eggplants that are to be eaten raw. Do not apply to eggplants that are to be processed. Do not apply to eggplants that are to be stored.

GRAPES: To control grape scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to grapes that are to be eaten raw. Do not apply to grapes that are to be processed. Do not apply to grapes that are to be stored.

CITRUS: (Lemon, lime, orange, tangerine, grapefruit) To control citrus scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to citrus that are to be eaten raw. Do not apply to citrus that are to be processed. Do not apply to citrus that are to be stored.

Directions for Use & Recommendations

To control Red Scale, Half Scale, Citrus Scale, Purple Scale, and other citrus scale insects, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to citrus that are to be eaten raw. Do not apply to citrus that are to be processed. Do not apply to citrus that are to be stored.

California and Arizona Only: Citrus Thrips 10-12 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to citrus that are to be eaten raw. Do not apply to citrus that are to be processed. Do not apply to citrus that are to be stored.

Do not apply more than 2000 gal. spray per acre. Do not apply more than once per season on lemons and oranges. Do not apply within 21 days of harvest, on other citrus fruits. Do not apply within 90 days of previous application. On tangerines, do not apply more than twice per season. Do not apply oil within 3 weeks of a sulfur application. Do not use oil on navel oranges.

MELON: To control melon scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to melons that are to be eaten raw. Do not apply to melons that are to be processed. Do not apply to melons that are to be stored.

ONIONS: To control onion maggot, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to onions that are to be eaten raw. Do not apply to onions that are to be processed. Do not apply to onions that are to be stored.

PEACHES, NECTARINES: European Red Mite, 1.0 lb. plus 1 lb. of sulfur per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to peaches and nectarines that are to be eaten raw. Do not apply to peaches and nectarines that are to be processed. Do not apply to peaches and nectarines that are to be stored.

NECTARINES: To control nectarine scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to nectarines that are to be eaten raw. Do not apply to nectarines that are to be processed. Do not apply to nectarines that are to be stored.

PEPPERS, PIMENTOS: To control pepper scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to peppers and pimentos that are to be eaten raw. Do not apply to peppers and pimentos that are to be processed. Do not apply to peppers and pimentos that are to be stored.

PLUMS, PRUNES: European Red Mite, 1.0 lb. plus 1 lb. of sulfur per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to plums and prunes that are to be eaten raw. Do not apply to plums and prunes that are to be processed. Do not apply to plums and prunes that are to be stored.

STRAWBERRY: To control strawberry scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to strawberries that are to be eaten raw. Do not apply to strawberries that are to be processed. Do not apply to strawberries that are to be stored.

TOMATOES: To control tomato scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to tomatoes that are to be eaten raw. Do not apply to tomatoes that are to be processed. Do not apply to tomatoes that are to be stored.

WALNUT: To control walnut scale, apply 1.5 lb. per acre at the start of the growing season. Do not apply within 21 days of harvest. Do not apply to walnuts that are to be eaten raw. Do not apply to walnuts that are to be processed. Do not apply to walnuts that are to be stored.



ETHION 25-WP

**ORGANOPHOSPHORUS INSECTICIDE-ACARICIDE
WETTABLE POWDER—FOR USE IN WATER**

ACTIVE INGREDIENT:

Ethion (O,O,O',O'-tetraethyl S,S'-methylene bisphosphorodithioate) 25.0%

INERT INGREDIENTS:

75.0%

100.0%

WARNING

Keep Out of
Reach of Children

**MAY BE FATAL IF SWALLOWED.
ABSORBED THROUGH SKIN.**

DO NOT BREATHE VAPOR, DUST OR SPRAY MIST.
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
WASH THOROUGHLY AFTER HANDLING.
DO NOT CONTAMINATE FOOD OR FEED.

476-1792
ACCEPTED
Jan 16 1971

USE PRECAUTIONS

BURN EMPTY BAG KEEP OUT OF SMOKE

LB. NET

STAUFFER CHEMICALS