

ACCEPTED

APR 29 1967

UNDER THE FEDERAL INSECTICIDE ACT AND REGULATIONS THEREUNDER FOR ECONOMIC POISON REGISTERED UNDER NO. 476-1798

CRYOLITE SEVIN 35-5 DUST

AN AGRICULTURAL INSECTICIDE

ACTIVE INGREDIENTS:

Sodium Fluoaluminate 35%
Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS 60%
100%

¹Trade-Mark of Union Carbide Corp. for the active ingredient, 1-naphthyl N-methylcarbamate.

CAUTION ✓

HARMFUL IF INHALED OR SWALLOWED

Do not take internally. Avoid breathing dust. Avoid skin contact.

WASH THOROUGHLY AFTER HANDLING.

Wash hands before eating and bathe after work. Wear long sleeved clothing. Change to clean clothing daily.

✓ **Keep Out of Reach of Children**

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Do not apply during bloom when honey bees or other pollinators are active. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after pollination.

To avoid pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or domestic animals or into adjoining food, fiber or pasture crops. The grower is responsible for residues on his property as well as for damages caused by drift from his property to that of others.

Consult state agricultural extension service or state agricultural experiment stations for information as the timing, number, and rate of applications needed under local conditions.

KEEP IN EMPTY BAG - KEEP OUT OF SMOKE ✓

50 LB. NET

^{power or} DIRECTIONS FOR USE

Apply when air is calm, using ground equipment, aircraft, ~~hand held~~ make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe the limits given.

RECOMMENDATION

LETTUCE: Armyworms, cabbage caterpillars, climbing cutworms, corn earworm, flea beetles, harlequin bugs, leafhoppers, lygus bugs, striped leafhoppers, stink bugs. Apply 25-40 lbs. per acre at first sign of infestation and repeat when necessary. Do not apply after seedling stage.

DO NOT STORE NEAR FOOD OR FEED.

NOTICE: Stauffer Chemical Company makes no warranty of merchantability or any other express or implied warranty concerning this material. It shall not be held responsible for personal injury, property damage, or other loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability resulting from such handling, storage or use.

UNSA Reg. No. 476-1798

Stauffer Chemical Company
P.O. Box 1000, Des Moines, Iowa 50319

Handwritten:
3-21-67