



# CAPTAN 25 SEED PROTECTANT

**PLANTER BOX OR DRILL BOX APPLICATION  
FOR SEED DISEASE CONTROL  
SPECIAL LUBRICANT ADDED — DISCOLORED**

**ACTIVE INGREDIENTS:**

CAPTAN: N-(trichloromethyl)thio-1,4-cyclohexene-1,2-dicarboximide..... 23.64%

Related derivatives..... 1.36%

**INERT INGREDIENTS:**

75.00%

100.00%

## CAUTION

**Keep Out of Reach of Children**

**HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.**

Avoid inhaling dust — wear respirator. Avoid contact with skin, eyes or clothing. Wash with soap and water after handling. Do not contaminate feed or food.

**PROTECT FROM EXCESSIVE HEAT — STORE IN A COOL DRY PLACE.**

**NOTICE — READ CAREFULLY**

**CONDITIONS OF SALE:** Stauffer (and seller) offer(s) this product for sale subject to, and buyer and all users are deemed to have accepted, the following conditions of sale and warranty which may only be waived by written agreement of a duly authorized representative of Stauffer.

**WARRANTY LIMITATION:** Stauffer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use on the label subject to the inherent risks referred to below. Stauffer makes no other express warranties; THERE IS NO IMPLIED WARRANTY OF MERCHANTABILITY and there are no warranties which extend beyond the description on the label hereof.

**INHERENT RISKS:** The directions for use of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use or application of this product contrary to label instructions or resulting from extra-ordinary weather conditions.

**LIMITATION OF LIABILITY:** In no case shall Stauffer be liable for special, indirect, or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

This product is sold only for uses stated on the label. No express or implied license is granted to use or sell this product under any patent in any country except as specified: Country, U.S.A.

Made In U.S.A. By

**STAUFFER CHEMICAL COMPANY**

**WESTPORT, CT 06880**

### PRODUCT INFORMATION

CAPTAN 25 SEED PROTECTANT is especially formulated to protect seed from seed-borne and soil-borne fungi. **PLANT SAFETY:** At the recommended dosage CAPTAN causes no reduction. The germination of treated seed is the same as from other seed fungicides.

### GENERAL USE PRECAUTIONS

Read all label directions before using and use only as directed. **IMPORTANT:** Treated seed must not be used for feed, food or oil purposes.

**FISH CAUTIONS:** Toxic to fish. Keep out of lakes, streams or disposal of wastes. Dispose of excess treated seed.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product on any crop not listed on the label. The amount of seed protectant, as shown in the directions, should be applied to the seed in the drill box and stirred with the seed for uniform coverage of the seed and any soil in the furrow, giving additional protection to the treated seed.

### RECOMMENDATIONS

**CORN (Field):** Use at the rate of 4 ounces per bushel or 4 to 6 ounces per 100 pounds of seed.

**PEANUTS (Shelled):** For control of Aspergillus use at the rate of 4 to 6 ounces per 100 pounds of seed.

**RICE:** Use at the rate of 6 ounces per bushel or 6 to 8 ounces per 100 pounds of seed.

**SOYBEANS:** Use at the rate of 4 ounces per bushel or 4 to 6 ounces per 100 pounds of seed.

**WHEAT:** Use at the rate of 3 ounces per bushel or 3 to 4 ounces per 100 pounds of seed. Treatment is not recommended for control of Blight.

**STORAGE:** Store in a cool, dry place. Do not store in areas where food is stored.

**PROHIBITIONS:** Do not contaminate water, food, or feed with this product.

**PESTICIDE DISPOSAL:** Pesticide, or rinsate that cannot be used, should be disposed of in a safe place as approved for pesticides or buried in a safe place as approved for pesticides.

**CONTAINER DISPOSAL:** Dispose of in an incinerator or by other means approved for pesticides.

**GENERAL:** Consult federal, state or local disposal regulations.

EPA Reg. No. 476-1808 EPA Est. No. 1-1000

# 25 SEED PROTECTANT

**DRILL BOX APPLICATION  
CONTROL  
D - DISCOLORED**

.....	23.64%
.....	1.36%
.....	75.00%
.....	100.00%

**Keep Out of Reach of Children**

**AVOID CONTACT WITH SKIN.**  
Wash with soap and water after use of clothing.

**STORE IN A COOL DRY PLACE.**

Buyer and all users are deemed to have accepted, the following authorized representative of Stauffer.  
Information on the label and is reasonably fit for the purposes referred to. Stauffer makes no other express warranties; THERE IS NO IMPLIED warranty on the label hereof.  
Should be followed carefully. However, it is impossible to eliminate all damages resulting from the use or handling of this product in respect of which such damages are claimed.  
No express or implied license is granted to use this product in any country other than that specified: Country: U.S.A.

**STAUFFER COMPANY**

6880

### PRODUCT INFORMATION

CAPTAN 25 SEED PROTECTANT is especially formulated for the treatment of seed in the planter box or drill box to protect seed from seed-borne and soil-borne fungi which cause Damping Off, Seed Decay and Seedling Blight.

**PLANT SAFETY:** At the recommended dosage CAPTAN 25 is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated giving a superior result as compared to results from other seed fungicides.

### GENERAL USE PRECAUTIONS

Read all label directions before using and use only as specified on this label.

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Do not use treated seed for feed, food or oil purposes.

**FISH CAUTIONS:** Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Dispose of excess treated seeds by burying with wastes.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

The amount of seed protectant, as shown in the recommendations, should be poured over the seed in the planter or drill box and stirred with the seed for uniform distribution throughout the hopper. The amounts recommended will provide adequate coverage of the seed and any excess, that does not adhere to the seed coat, will flow with the seed into the furrow, giving additional protection against soil-borne diseases. This product may be used on previously treated seed.

### RECOMMENDATIONS

**CORN (Field):** Use at the rate of 4 ounces per bushel or 7 ounces per 100 pounds of seed as a planter box treatment.

**PEANUTS (Shelled):** For control of Aspergillus Niger and other seed and soil-borne diseases, use 2 to 4 ounces per bushel or 4 to 6 ounces per 100 pounds of seed as a planter box treatment.

**RICE:** Use at the rate of 6 ounces per bushel or 14 ounces per 100 pounds of seed as a planter box treatment.

**SOYBEANS:** Use at the rate of 4 ounces per bushel or 7 ounces per 100 pounds of seed as a planter box treatment.

**WHEAT:** Use at the rate of 3 ounces per bushel or 4½ ounces per 100 pounds of seed as a drill box treatment. This treatment is not recommended for control of Bunt (Smut).

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE DISPOSAL:** Pesticide, or rinseate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

**CONTAINER DISPOSAL:** Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

**GENERAL:** Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

EPA Reg. No. 476-1808

EPA Est. No. 476-

10 LBS. NET



5

6



# CAPTAN 25 SEED PROTECTANT

**PLANTER BOX OR DRILL BOX APPLICATION  
FOR SEED DISEASE CONTROL  
SPECIAL LUBRICANT ADDED — DISCOLORED**

**ACTIVE INGREDIENT:**

Captan\*

**INERT INGREDIENTS:** .....

\*N-[(trichloromethyl)thio]-4-cyclohexene-1,2,-dicarboximide.

## CAUTION

**Keep Out of Reach of Children**

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN.

Avoid inhaling dust — wear respirator. Avoid contact with skin, eyes or clothing. Wash with soap and water after handling. Do not contaminate feed or food.

PROTECT FROM EXCESSIVE HEAT — STORE IN A COOL DRY PLACE.

CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURSTING WITH WATER.

EPA REG. NO. 476-1808

**10 LBS. NET**


Made In U.S.A. By

*June 19, 1993*

**STAUFFER CHEMICAL COMPANY**

**WESTPORT, CT 06880**

*476-1808*



# CAPTAN 25 SEED PROTECTANT

## OR DRILL BOX APPLICATION

### ED DISEASE CONTROL

#### ICANT ADDED — DISCOLORED

25%  
75%  
100%

carboximide.

**Keep Out of Reach of Children**

ABSORBED THROUGH SKIN.  
Avoid contact with skin, eyes or clothing. Wash with soap and water after

AVOID EXCESSIVE HEAT — STORE IN A COOL DRY PLACE.  
DISPOSED BY CRUSHING AND BURYING WITH WASTES.

10 LBS. NET

A-1

Made in U.S.A. By

**CHEMICAL COMPANY**  
HARTFORD, CT 06180

### PRODUCT INFORMATION

CAPTAN 25 SEED PROTECTANT is especially formulated for the treatment of seed in the planter box or drill box to protect seed from seed-borne and soil-borne fungi which cause Damping Off, Seed Decay and Seedling Blight.

**PLANT SAFETY:** At the recommended dosage CAPTAN 25 is non-injurious from the standpoint of seed injury or yield reduction. The germination of treated seed is frequently stimulated giving a superior result as compared to results from other seed fungicides.

### GENERAL USE PRECAUTIONS

Read all label directions before using and use only as specified on this label.

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Do not use treated seed for feed, food or oil purposes.

**FISH CAUTIONS:** Toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Dispose of excess treated seeds by burying with wastes.

### DIRECTIONS FOR USE

The amount of seed protectant, as shown in the recommendations, should be poured over the seed in the planter or drill box and stirred with the seed for uniform distribution throughout the hopper. The amounts recommended will provide adequate coverage of the seed and any excess, that does not adhere to the seed coat, will flow with the seed into the furrow, giving additional protection against soil-borne diseases. This product may be used on previously treated seed.

### RECOMMENDATIONS

**CORN (Field):** Use at the rate of 4 ounces per bushel or 7 ounces per 100 pounds of seed as a planter box treatment.

**PEANUTS (Shelled):** For control of Aspergillus Niger and other seed and soil-borne diseases, use 2 to 4 ounces per bushel or 4 to 6 ounces per 100 pounds of seed as a planter box treatment. *damping off, mold, seedling blight*

**RICE:** Use at the rate of 6 ounces per bushel or 14 ounces per 100 pounds of seed as a planter box treatment.

**SOYBEANS:** Use at the rate of 4 ounces per bushel or 7 ounces per 100 pounds of seed as a planter box treatment.

**WHEAT:** Use at the rate of 3 ounces per bushel or 4½ ounces per 100 pounds of seed as a drill box treatment. This treatment is not recommended for control of Bunt (Smut).

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.