

WEEDS CONTROLLED

Smooth Crabgrass	(Digitaria ischaemum)
Hairy Crabgrass	(Digitaria sanguinalis)
Redroot Pigweed	(Amaranthus retroflexus)
Watergrass	(Echinochloa spp.)
Lambsquarters	(Chenopodium album)
Goosegrass	(Eleusine indica)
Annual Bluegrass	(Poa annua)
Shepherdspurse	(Capsella bursa-pastoris)
Deadnettle	(Lamium amplexicaule)

GENERAL USE PRECAUTIONS

BETASAN should be used only for recommended uses and at recommended rates. Apply when there is little or no wind to avoid drift. Clean up spills on walks or patios by covering with dry soil or absorptive material. Bury sweepings and hose area. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Dispose of container by punching holes in it and burying with wastes. Do not allow children or pets on treated lawns until BETASAN has been watered into soil and grass is completely dry. Avoid contamination of food and feed. Toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.

PRODUCT INFORMATION

BETASAN is recommended for use on home lawns, parks and golf courses, on ornamental plants and on ground covers for control of crabgrass, annual bluegrass and other annual weeds listed.

DIRECTIONS FOR APPLICATION

Application of BETASAN must precede emergence of weeds from the soil. Crabgrass will emerge from the soil from four to six weeks before it is visible above lawn sod. If weeds are visible above the soil, it is too late to apply BETASAN. Remove leaves, dead or tall grass, and other debris, or thoroughly work them into the soil before applying BETASAN.

TIMING OF APPLICATION

GENERAL COMMENTS

GRASS LAWNS: BETASAN should be applied only to well-established grass lawns. BETASAN can be used on the following established turf grasses:

Bahia	Centipede	Poa Trivialis
Bentgrass spp.	Fescue spp.	St. Augustine
Bermudagrass	Pensacola	Red Top
Perennial Bluegrass spp.	Perennial Ryegrass	Zoysia

DICHONDRA LAWNS: BETASAN can be applied to dichondra lawns at the time of seeding or any time thereafter.

ORNAMENTALS AND GROUND COVERS: BETASAN should be applied only to the following ornamentals and ground covers.

Alyssum	Coral Bell	Stock
Aster	Daisy	Sweet Pea
Bachelor's-Button	Marigold	Wall Flower
Calendula	Pansy	Zinnia
Candy-Tuft	Primrose	
Daffodil	Gladiolus	Ranunculus
Dahlia	Narcissus	Tulip
Freesia		
Abelia	Juniper	Pyracantha
Azalea	Monterey Cypress	Sandankwa
Azara	Monterey Pine	Star Jasmine
Boxwood	Myrtle	Tobira
Daphne	Privet	Xylosma
Holly		
Ajuga	Hypericum	Periwinkle
Campanula	Ice Plant	Sedum
Festuca	Ivy	Wild Strawberry
Gazania	Pachysandra	

Betasan[®] 4-E

1 GAL. NET

A SELECTIVE HERBICIDE FOR PRE-EMERGENCE CONTROL OF CRABGRASS AND ANNUAL BLUEGRASS IN TURF, ORNAMENTALS AND GROUND COVERS

Contains 4 lb. BETASAN per Gallon

ACTIVE INGREDIENT:

S-(O, O-diisopropyl phosphorodithioate) of N-(2-mercaptoethyl) benzenesulfonamide

45.2%

INERT INGREDIENTS

54.8%

®Registered T.M. of Stauffer Chemical Company — U.S. Patent No. 3,205,253

100.0%

CAUTION

- HARMFUL IF SWALLOWED.
- AVOID PROLONGED OR REPEATED CONTACT WITH THE SKIN.
- AVOID CONTACT WITH THE EYES.
- AVOID CONTAMINATION OF FOOD AND FEED, DRINKING WATER, FISH PONDS, ETC.

Keep Out of Reach of Children

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME

READ ALL DIRECTIONS BEFORE USING.

STAUFFER CHEMICAL CO.

NEW YORK

ACCEPTED
 Mar 9, 1967
 FEDERAL AGENCY FOR ENVIRONMENTAL PROTECTION
 DIVISION OF PESTICIDES AND ROENTIGENIC AGENTS
 POISON REGISTER
 476-1817

476-1817

03/09/1967

1/2

:T

For control of CRABGRASS, apply BETASAN at the recommended rate (See "RATES OF APPLICATION") in the late winter or early spring, depending on the location and seasonal conditions.

For control of ANNUAL BLUEGRASS AND OTHER ANNUAL WEEDS, apply BETASAN at the recommended rate (See "RATES OF APPLICATION") before the weeds emerge from the soil. Consult State Agricultural Experiment Station Weed Specialists or Garden Weed Specialists for specific information as the timing of application will vary with local conditions.

In southern or western states where germination of crabgrass, annual bluegrass and other annual weeds extends into the late summer and early fall, a second application of BETASAN at the recommended rates should be made 4 to 5 months after the initial application.

BETASAN should be applied with spray equipment that will provide uniform application such as knapsack sprayer, hose proportioner, or pressure sprayer.

Sprinkle with water for 10 to 15 minutes after application to wash the BETASAN off the grass and down into the soil where the weed seeds are located.

RATES OF APPLICATION

CRABGRASS: Apply 1 gallon of BETASAN 4-E in 40 to 50 gallons of water uniformly on 17,400 square feet.
ANNUAL BLUEGRASS AND OTHER ANNUAL WEEDS: Apply 1 gallon of BETASAN 4-E in 40 to 50 gallons of water uniformly on 13,900 square feet.

RESEEDING

Any material that provides effective control of annual grasses may affect new lawn grass seedlings which germinate after treatment. Proper fertilization and watering will encourage the existing turf grass to fill in the thin or bare spots formerly occupied by crabgrass.

The following procedures are suggested in order to minimize any stunting of turf grasses which may occur when reseeding into BETASAN treated areas:

1. Do not reseed within four months after application.
2. Renovate the area to be reseeded by raking thoroughly and mix sterilized (weed-free) organic mulch (50-50 by volume) into the top two inches of soil.

NOTICE: *Stauffer Chemical Company makes no warranty of merchantability or any other express or implied warranty concerning this material.* It shall not be held responsible for personal injury, property damage, or other loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability resulting from such handling, storage or use.

USDA Reg. No. 476-1817

T ARC 660921 SE

Made in U.S.A

Stauffer Chemical Co., New York