

Stauffer
CHEMICALS

CAPTAN 80-WP

A Fungicide For Plant Disease Control

ACTIVE INGREDIENTS	
CAPTAN: N-[[trichloromethyl] thio]-4-cyclohexene-1,2-dicarboximide	75.7%
Related derivatives	4.3%
INERT INGREDIENTS	20.0%
	100.0%

CAPTAN 80-WP is a microfine wettable powder for use in water as a spray for the control of certain fungus diseases of fruit, vegetables, and ornamental crops, and as a soil treatment for the control of certain seed rots and damping-off diseases.

EPA Reg. No. 476-1819

CAUTION

Keep Out of Reach of Children

CAUSES SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED OR INHALED. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Use with adequate ventilation. Wash thoroughly after handling. Do not store near food or feed.

FIRST AID: In case of eye contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician. Wash skin with soap and water.

STORE IN COOL, DRY PLACE—PROTECT FROM EXCESSIVE HEAT

5 LB. NET WT. 2.27 kg

ACCEPTED

SEP 4 1981

STAUFFER CHEMICAL COMPANY
WESTPORT, CT 06881 U.S.A.

POST HARVEST DIP

Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

PLUMS PRUNES Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

QUINCE Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

STRAWBERRIES Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

VEGETABLE CROPS

ASPARAGUS (Preplanting Root Dip) Dip roots in solution before planting. Use 1/2 to 1 lb. per 100 lbs. of roots.

BEANS PEAS Apply to seed surfaces before planting. Use 1/2 to 1 lb. per 100 lbs. of seed.

CANTALOUPE CUCUMBER PUMPKIN SUMMER AND WINTER SQUASH HONEYDEW MELON WATERMELON Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

CARROTS Apply to root surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of root.

CELERY Apply to leaf surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of leaf.

EGGPLANT Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

ONIONS (GREEN) Apply to leaf surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of leaf.

ONIONS (Green and Bulb) Apply to bulb surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of bulb.

PEPPERS PIMENTOS Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

TOMATO Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

SPECIAL USES

PREPLANTING SOIL TREATMENT Apply to soil before planting. Use 1/2 to 1 lb. per 100 lbs. of soil.

BEANS PEAS Apply to seed surfaces before planting. Use 1/2 to 1 lb. per 100 lbs. of seed.


BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER COLLARDS KALE KOHLRABI MUSTARD RUTABAGAS TURNIPS Apply to leaf surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of leaf.

CANTALOUPE CUCUMBERS HONEYDEW MELONS PUMPKIN SQUASH (SUMMER AND WINTER) WATERMELONS Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

SPINACH Apply to leaf surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of leaf.

POST HARVEST DIP Apply to fruit surfaces immediately after harvest. Use 1/2 to 1 lb. per 100 lbs. of fruit.

Apples, Apricots, Cherries, Citrus, Grapes, Lemons, Limes, Mangoes, Nectarines, Onions (Green and Bulb), Oranges, Peaches, Pears, Potatoes, Tangerines, Tangelos.



11/2 2/10/11

Post Harvest Dip
PLUMS PRUNES
QUINCE
STRAWBERRIES

VEGETABLE CROPS
ASPARAGUS (Preplanting Root Dip)
BEANS PEAS

CANTALOUPE CUCUMBER PUMPKIN SUMMER AND WINTER
JASH. HONEYDEW MELON WATERMELON
CARROTS

CELERY
EGGPLANT
ONIONS (Green and Bulb)

PEPPERS MIMENTOS
TOMATO

SPECIAL USES
PREPLANTING SOIL TREATMENT

BEANS PEAS
BEETS
BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER
COLLARDS KALE KOHLRABI MUSTARD RUTABAGAS TURNIPS
CANTALOUPE CUCUMBERS HONEYDEW MELONS PUMPKIN
SQUASH (SUMMER AND WINTER) WATERMELONS

SPINACH
POST HARVEST DIP
Apples, Apricots, Cantaloupes, Cherries, Cucumbers, Grapefruit, Lemons, Limes, Mangoes, Nectarines, Onions (Green and Bulb), Oranges, Peaches, Pears, Potatoes, Tangelos, Tangerines

Stauffer
CHEMICALS

CAPTAN 80-WP

A Fungicide For Plant Disease Control

ACTIVE INGREDIENTS:	
CAPTAN, N-[(trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide	75.7%
Related derivatives	4.3%
INERT INGREDIENTS	20.0%
	100.0%

Captan 80-WP is a microfine wettable powder for use in water as a spray for the control of certain fungus diseases of fruit, vegetables, and ornamental crops, and as a soil treatment for the control of certain seed rots and damping-off diseases.

EPA Reg. No. 476-1819

CAUTION Keep Out of Reach of Children
CAUSES SKIN AND EYE IRRITATION. MAY BE HARMFUL IF SWALLOWED OR INHALED. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Use with adequate ventilation. Wash thoroughly after handling. Do not store near food or feed.

FIRST AID: In case of eye contact, immediately flush eyes with plenty of water for at least 15 minutes. Call a physician. Wash skin with soap and water.

STORE IN COOL, DRY PLACE—PROTECT FROM EXCESSIVE HEAT

5 LB. NET WT. 2.27 kg

MADE IN U.S.A. BY
STAUFFER CHEMICAL COMPANY
WESTPORT, CT 06881





CAPTAN 80-WP

After application, the active ingredient, captan, remains on the surface of the plant and is not absorbed. Captan is a contact fungicide and does not move through the plant. Captan is effective against a wide range of fungi and is used to control a wide range of diseases. Captan is a contact fungicide and does not move through the plant. Captan is effective against a wide range of fungi and is used to control a wide range of diseases.

CAUTION

PACKING BOX TREATMENT

For treatment of packing boxes, use 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Apply to the inside of the box and to the outside of the box. Repeat application every 7 days for 3 applications.

ORNAMENTALS

AZALEAS - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

BEGONIAS (TUBEROUS) - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

CAMELLIAS - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

CARNATIONS - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

CHRYSANTHEMUM - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

DICHONDRA - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

GLADIOLUS (CORMS) - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

GRASSES (ORNAMENTAL IN NON-PASTURED AREAS ONLY) - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

GRASSES (LAWN SEEDBEDS) - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

ROSES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

SOIL AND GREENHOUSE BENCH TREATMENT - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Read the entire label before using this product. Do not use this product if the label directions do not apply to the crop and conditions of use. Do not use this product if the label directions do not apply to the crop and conditions of use.

FISH CAUTIONS

This product is highly toxic to fish. Do not use this product in or near water. Do not use this product in or near water.

COMPATIBILITY AND PLANT SAFETY - Captan 80-WP is a contact fungicide and does not move through the plant. Captan 80-WP is a contact fungicide and does not move through the plant.

DIRECTIONS FOR USE

Use this product in a manner consistent with its labeling. Use this product in a manner consistent with its labeling.

RECOMMENDATIONS

Use this product in a manner consistent with its labeling. Use this product in a manner consistent with its labeling.

FRUIT AND NUT CROPS

IMPORTANT - Read the label carefully. Do not use this product if the label directions do not apply to the crop and conditions of use. Do not use this product if the label directions do not apply to the crop and conditions of use.

ALMONDS - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

APPLES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

APRICOTS - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

RASPBERRIES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

BLUEBERRIES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

CHERRIES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines) - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

CRANBERRIES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

GRAPES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

MANGOES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

PEACHES, NECTARINES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

PEACHES (Nursery Stock) - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

PEARS - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

PINEAPPLES - Control leaf spot, scale, and other diseases. Apply 1/2 lb. of Captan 80-WP per 100 sq. ft. of surface area. Repeat application every 7 days for 3 applications.

STORAGE AND DISPOSAL

PROHIBITIONS - Do not use this product in or near water. Do not use this product in or near water.

PESTICIDE DISPOSAL - Do not use this product in or near water. Do not use this product in or near water.

CONTAINER DISPOSAL - Do not use this product in or near water. Do not use this product in or near water.

GENERAL - Do not use this product in or near water. Do not use this product in or near water.

NOTICE—READ CAREFULLY

CONDITIONS OF SALE - This product is sold as a fungicide. This product is sold as a fungicide.

WARRANTY LIMITATION - This product is sold as a fungicide. This product is sold as a fungicide.

INHERENT RISKS - This product is sold as a fungicide. This product is sold as a fungicide.

LIMITATION OF LIABILITY - This product is sold as a fungicide. This product is sold as a fungicide.

STAUFFER CHEMICAL COMPANY





Page 2 of 2 sections

CAPTAN 80-WP



CAUTION - See the following use precautions for information on the hazards of this product.

PACKING BOX TREATMENT - To protect the contents of the packing box from moisture damage, the packing box should be treated with Captan 80-WP. Apply 1/2 ounce of Captan 80-WP per 100 square feet of packing box surface. Repeat application every 14 days.

ORNAMENTALS
AZALEAS - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.
BEGONIAS (TUBEROUS) - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

CAMELIAS - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.
CARNATIONS - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

ADIOLUS (CORMS) - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.
GRASSES (ORNAMENTAL IN NON-PASTURED AREAS ONLY) - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

GRASSES (LAWN SEEDBEDS) - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.
ROSES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

SOIL AND GREENHOUSE BENCH TREATMENT - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.
CHRYSANTHEMUM - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

STORAGE AND DISPOSAL
PROHIBITIONS - Do not store in areas where food or feed is produced, stored, or consumed.
PESTICIDE DISPOSAL - Pesticide should be disposed of in accordance with local, state, and federal regulations.
CONTAINER DISPOSAL - Containers should be disposed of in accordance with local, state, and federal regulations.
GENERAL - Store in a cool, dry place. Do not use if the container is damaged or leaking.

NOTICE—READ CAREFULLY
CONDITIONS OF SALE - This product is sold as a fungicide. It is not a guarantee of performance. Use according to label directions.

WARRANTY LIMITATION - The manufacturer warrants that the product is free from defects in material and workmanship at the time of sale. No other warranties are made.

INHERENT RISKS - The use of this product may result in the death of the plant or animal to which it is applied. Use with caution.

LIMITATION OF LIABILITY - The manufacturer shall not be liable for any damages, including consequential damages, arising from the use of this product.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Read the directions and precautions for use on the label. See the following use precautions for information on the hazards of this product.

FISH CAUTIONS - Do not use in or near water. Do not use in or near streams, rivers, lakes, or ponds. Do not use in or near fish hatcheries or other aquatic areas.

COMPATIBILITY AND PLANT SAFETY - Captan 80-WP may be incompatible with other fungicides. Do not use in combination with other fungicides unless the label directions are followed. Captan 80-WP may be phytotoxic to certain plants. Do not use on plants listed in the "Plant Safety" section of the label.

DIRECTIONS FOR USE

Apply Captan 80-WP in the manner specified on the label. Do not use in or near water. Do not use in or near streams, rivers, lakes, or ponds.

RECOMMENDATIONS - Use the maximum permissible rate per acre per application. Do not use in or near water. Do not use in or near streams, rivers, lakes, or ponds.

FRUIT AND NUT CROPS

IMPORTANT - Read the label directions for use on fruit and nut crops. Do not use in or near water. Do not use in or near streams, rivers, lakes, or ponds.

ALMONDS - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

APPLES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

CAUTION - Do not use in or near water. Do not use in or near streams, rivers, lakes, or ponds.

APRICOTS - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

BLACKBERRY, DEWBERRY, RASPBERRY - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

CHERRIES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangelos) - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

GRAPES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

MANGOES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

PEACHES, NECTARINES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

PEARS - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

PINEAPPLES - Apply 1/2 ounce of Captan 80-WP per 100 square feet of foliage. Repeat application every 14 days.

STAUFFER CHEMICAL COMPANY

