

ACCEPTED
2-4-70
476-1830

CAPTAN- SEVIN* 5-2.5 DUST

AGRICULTURAL INSECTICIDE AND FUNGICIDE

ACTIVE INGREDIENTS:

CAPTAN: N[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide	5.0%
Carbaryl (1-naphthyl N-methylcarbamate)	2.5%
INERT INGREDIENTS	92.5%

Not licensed for use in countries where patents still exist.

*Registered trademark of Union Carbide Corporation for the active ingredient (carbaryl).

CAUTION

Keep Out of Reach
of Children

HARMFUL IF INHALED OR SWALLOWED.

Avoid prolonged or repeated breathing of dust.
 Avoid prolonged or repeated contact with skin.
 Do not take internally.
 Wash hands and face before eating.
 Take shower or bath after work.
 Wear regular long sleeved work clothing. Change to clean clothing daily.
 Avoid contamination of food, feed, water supplies, streams and ponds.
NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

STORE IN A COOL DRY PLACE; PROTECT FROM EXCESSIVE HEAT.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

Made in U.S.A. — Stauffer Chemical Co., New York

USDA Reg. No. 476-1830

S-SF-691204-NP

USE PRECAUTIONS

Use only for claims listed and in accordance with all label directions. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of dust from areas treated.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Consult state agricultural extension service or state agricultural experiment stations for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

DIRECTIONS FOR USE

Sevin insecticide will not control insects if used with lime or other highly alkaline materials. Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7-10 day intervals when needed to maintain control.

RECOMMENDATIONS (ALSO READ USE PRECAUTIONS & DIRECTIONS FOR USE)

STRAWBERRIES: Botrytis fruit rot (gray mold), leaf spot, meadow spittlebug, strawberry leaf roller, strawberry weevil.
Apply 40-50 lb. per acre. May be applied up to one day before harvest.

BURN EMPTY BAG — KEEP OUT OF SMOKE. 50 LB. NET