



CAPTAN SP

AGRICULTURAL FUNGICIDE AQUEOUS SUSPENSION 4 FLOWABLE SEED PROTECTANT

ACTIVE INGREDIENT:
CAPTAN N-(trichloromethyl)thio-4-cyclohexene-1,
2-dicarboximide

100%
61.5

100.0%

(Contains 4 lb. Captan Per Gallon)

CAUTION Keep Out of Reach of Children
HARMFUL IF TAKEN INTERNALLY
Avoid inhalation of spray.
Avoid prolonged or repeated contact with skin.
Avoid contamination of feed and feed.

PRODUCT FREEZING TEMPERATURE 32 F

DO NOT REUSE CONTAINER. WASH THOROUGHLY AND DISCARD IN A SAFE PLACE

1 GAL NET

STAUFFER CHEMICAL COMPANY
NEW YORK, NY 10017

SEED TREATMENT DOSAGE RATES

Seed	Dosage
Alfalfa, Clover, Lespedeza, Trefol...	8 1/4 fl. oz. 100 lb
Beans (snap), dry lima, Cowpeas, Peas...	2 1/2 fl. oz. 100 lb
Beets (garden)	12 fl. oz. 100 lb
Blue Grass	8 1/4 fl. oz. 100 lb
Cereal Grains (for seed and seedling rats - will not control bunt smut)	
Barley	2 1/2 fl. oz. 100 lb
Oats	3 1/2 fl. oz. 100 lb
Wheat	1 1/2 fl. oz. 100 lb
Cole Crops (broccoli, Brussels sprouts, cabbage, cauliflower)	1 1/2 fl. oz. 100 lb
Field Peas	1 1/4 fl. oz. 100 lb
Shelf Beans	1 1/2 fl. oz. 100 lb
Wild delinited	3 fl. oz. 100 lb
Red clover	3 fl. oz. 100 lb
Fuzzy	4 1/2 fl. oz. 100 lb
Winter cantaloupe	3 fl. oz. 100 lb
Corn Meal	1 1/2 fl. oz. 100 lb
Flax	3 1/2 fl. oz. 100 lb
Trinseeds	8 1/4 fl. oz. 100 lb
Milk	4 fl. oz. 100 lb
Onions (Planting)	1 1/2 pt/lb
Peanut	5 fl. oz. 100 lb
Pepper (California wonder)	3 fl. oz. 100 lb
Rice	1 fl. oz. 100 lb
Signon, manure, Ear kernels, oat	1 fl. oz. 100 lb
Soybean	2 1/2 fl. oz. 100 lb
Spinach	1 fl. oz. 100 lb
Squash, pumpkin, watermelon, muskmelon	1 fl. oz. 100 lb
Sweet Peas	1 fl. oz. 100 lb
Wheat	1 1/2 fl. oz. 100 lb
Yeast	1 1/2 fl. oz. 100 lb
Yard Chard	1 fl. oz. 100 lb

GRAPPLER PILES - Prevent leaf blight by spraying with 1 1/2 fl. oz. per 100 lb of fruit at maturity 20 days before harvest.