

SOUTHWESTERN STATES



READ ENTIRE LABEL BEFORE USING

Vernam 10 G

Contains 10 Pounds VERNAM per 100 Pounds

4-22-69 A SELECTIVE HERBICIDE
20/40 Mesh (U.S. Sieve Series Designation)
476-1872

50 LB. NET

ACTIVE INGREDIENT:

S-propyl dipropylthiocarbamate 10.0%

INERT INGREDIENTS

..... 90.0%

..... 100.0%

U. S. Pat. No. 2,913,327

CAUTION

Keep Out of Reach of Children

- Harmful if swallowed.
- Avoid contact with skin and clothing.
- Avoid contamination of food and feed.
- Wash thoroughly after use.
- Avoid breathing dust.

WEEDS CONTROLLED

ANNUAL GRASSES:

Barnyardgrass (Watergrass)
(Echinochloa spp.)

Crabgrass (Digitaria spp.)

Giant Foxtail (Setaria faberii)

Green Foxtail (Setaria viridis)

Yellow Foxtail (Setaria glauca)

German Millet (Setaria italica)

Goosegrass (Eleusine indica)

Wild Cane (Sorghum spp.)

Johnsongrass seedlings (Sorghum
halapense)

ANNUAL BROADLEAVED WEEDS:

Carpetweed (Molluga verticillata)

Sicklepod (Coffeeweed) Cassia
tora)

Florida Pusley (Richardia scabra)

Lambsquarters (Chenopodium
album)

Annual Morningglory (Ipomea
purpurea)

Pigweed (Smooth) (Amaranthus
hybridus)

Redroot Pigweed (Amaranthus
retroflexus)

Purslane (Portulaca oleracea)

Velvetleaf (Burrweed) (Abutilon
theophrasti)

The above listed broadleaf weeds will be controlled only if the application is made when conditions are favorable for weed seed germination and growth.

Perennial Weeds:

Yellow Nutgrass (Fenugreek)
Purple Nutgrass (Fenugreek)

Existing stands of nutgrass must be controlled by other means. The above listed weeds will be controlled only if the application is made when conditions are favorable for weed seed germination and growth.

USE PRECAUTIONS

Apply according to directions and label. Do not apply to growing crops. VERNAM will control weeds during germination and early stages of growth. Do not apply to crops during germination and early stages of growth.

Do not apply to crops during germination and early stages of growth. Do not apply to crops during germination and early stages of growth.

Do not apply to crops during germination and early stages of growth. Do not apply to crops during germination and early stages of growth.

Do not apply to crops during germination and early stages of growth. Do not apply to crops during germination and early stages of growth.

DIRECTIONS FOR USE OF VERNAM

VERNAM is a selective herbicide for use on broadleaf weeds in corn, sorghum, and other crops. It is applied as a pre-emergent or early post-emergent treatment. For best results, apply when weeds are beginning to germinate.

APPLICATION DIRECTIONS

Apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

Apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

INCORPORATION DIRECTIONS

INCORPORATION BEFORE PLANTING: Use a shallow tillage implement such as a harrow or disk to incorporate VERNAM into the soil to a depth of 4 to 6 inches (no deeper than 6 inches) followed by a 50% to 75% soil moisture.

INCORPORATION AFTER PLANTING: Use a shallow tillage implement such as a harrow or disk to incorporate VERNAM into the soil to a depth of 4 to 6 inches (no deeper than 6 inches) followed by a 50% to 75% soil moisture.

INCORPORATION AFTER PLANTING: Use a shallow tillage implement such as a harrow or disk to incorporate VERNAM into the soil to a depth of 4 to 6 inches (no deeper than 6 inches) followed by a 50% to 75% soil moisture.

PLANTING

For applications made before planting, use a shallow tillage implement such as a harrow or disk to incorporate VERNAM into the soil to a depth of 4 to 6 inches (no deeper than 6 inches) followed by a 50% to 75% soil moisture.

CULTURAL PRACTICES FOLLOWING APPLICATIONS

VERNAM is not persistent in the soil and susceptible weeds germinating later in the season may not be controlled. Cultivate to control these late germinating weeds, as well as the susceptible weeds that germinate before incorporation.

RECOMMENDATIONS

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

For best results, apply VERNAM to the soil in a band 4 to 6 inches wide around the base of the crop. Do not apply to the foliage of the crop. Do not apply to crops during germination and early stages of growth.

BURN EMPTY BAG—KEEP OUT OF SMOKE

STAUFFER CHEMICAL COMPANY

NEW YORK, NY 10017

NOTICE: STAUFFER CHEMICAL COMPANY MAKES NO WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY AND OR FITNESS FOR ANY PARTICULAR PURPOSE CONCERNING THIS MATERIAL, EXCEPT THOSE WHICH ARE CONTAINED ON STAUFFER'S LABEL.