

CAPTAN-PCNS 10-10 GRANULAR

SOIL FUNGICIDE SEED PROTECTANT

10/10 MEQR (U.S. Slave Series Designation)

ACTIVE INGREDIENTS:

CAPTAN-N-[(trichloromethyl) thio]-4-cyclohexene-2,1-dicarboximide	10%
Pentachloronitrobenzene	10%
INERT INGREDIENTS	80%
	100%

U. S. Pat. Nos. 2,775,776; 2,835,771; 2,555,775. Not licensed for use or export outside the United States, its territories, possessions, Canada or Mexico.

STAUFLER CHEMICAL COMPANY

Keep Out of Reach of Children

- MAY CAUSE SKIN IRRITATION.
- AVOID BREATHING DUST.
- AVOID CONTACT WITH SKIN AND EYES.
- WEAR GLOVES, GOGGLES AND RESPIRATOR WHEN LOADING EQUIPMENT.
- WASH HANDS AND FACE THOROUGHLY AFTER USING AND BEFORE EATING OR SMOKING

STAUFLER CHEMICAL COMPANY

Avoid drift to adjoining food and forage crops.

Toxic To Fish - Keep out of lakes, streams & ponds. 476-1881

NOTICE: Stautler Chemical Company makes no warranty, express or implied, including the merchantability and/or fitness for any particular purpose, of this material, except those which are contained on Stautler's label.

RECOMMENDATIONS

CAPTAN-PCNS 10-10 GRANULAR is a special formulation for use at time of planting to protect beans and cotton seed and seedlings against soil borne diseases.

DIRECTIONS FOR USE

COTTON

For protection against damping-off, seed rot, Rhizoctonia (Soruskin) and Phythium.

IN-FURROW APPLICATION: Apply 15 pounds of Captan-PCNS 10-10 Granular per acre based on a 30" row spacing. Increase the rate per acre proportionately as the row spacing decreases. For a 24" row spacing, use 18 1/2 pounds per acre; on an 18" row spacing, use 25 pounds per acre, and on a 14" row spacing, use 30 pounds per acre. Application should be made to the planting furrow and covering soil at the time of planting.

PLANTER BOX APPLICATION: Use 5 pounds of Captan-PCNS 10-10 Granular per acre. Mix seed and the proper amount of soil, moisture, fertilizer and insecticide in a container just prior to planting, and mix thoroughly. An equal amount of chemical may be added to the seed in the planter box at the time of sowing or mixing, where grain planter box application is not possible as a substitute for regular soil treatment. It is a substitute for a pre-plant chemical treatment of good seed.

WIND-DROP SEEDING — PLANTER BOX METHOD: Use 10-15 pounds of Captan-PCNS 10-10 Granular for wind-drip seeding, rate to be 10 pounds per acre. Balance of directions for planter box application. Wind-drip apply to wind-drip seedling method.

IN-FURROW APPLICATION: For protection against damping-off, seed rot, Rhizoctonia (Soruskin) and Phythium, apply 10-15 pounds of Captan-PCNS 10-10 Granular per acre based on a 30" row spacing. Increase the rate proportionately as the row spacing decreases. Application should be made to the planting furrow and covering soil at the time of planting.

NOTE: Do not use except planting seed and seedlings. Do not use for soil treatment. Do not use for seedling the seed and seedling. Do not use for seedling the seed and seedling. Do not use for seedling the seed and seedling.

S-57-507501-0-1-1