

ELECTRIC® TRITHION®

90-2 CITRUS DUST

ORGANIC PHOSPHATE INSECTICIDE-ACARICIDE

476-1481
ACC.
Jul 30 1968
U.S. DEPT. OF AGRICULTURE
FEDERAL INSECTICIDE AND FUNGICIDE ACT
REGISTERED TOXIC SUBSTANCE
EPA REG. NO. 476-1481

ACTIVE INGREDIENTS:

Sulfur	90%
Carbophenothion—S-[[[p-chlorophenyl]thio]methyl] O,O-diethyl phosphorodithioate	2%

INERT INGREDIENTS:

8%
100%

U. S. Pat. No. 2,793,224

FINENESS Not less than 93% passing 325 mesh

CAUTION

Keep Out of Reach of Children

HAZARDOUS IF SWALLOWED

Avoid breathing dust, contact with skin and eyes.

Wear dust respirator and goggles.

Wash hands, arms and face thoroughly with soap and water before eating or smoking.

Do not contaminate feed and food.

Wash all contaminated clothing with soap and hot water before re-use.

ANTIDOTE—TRITHION, upon repeated, prolonged or careless use, may cause cholinesterase inhibition. **Atropine is antidotal.**

USE PRECAUTIONS

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damages caused by drift from his property to that of others.

Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

Sulfur may cause severe fruit and leaf injury to sulfur-sensitive crops. Do not apply to cucurbits (cucumbers, cantaloupes, melons, squash) or other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90°F.

Do not use sulfur with oil sprays or within 30 days of an oil spray. Do not use sulfur on any crop unless local use has proved safe for plants.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control.

RECOMMENDATIONS

CITRUS: Citrus red (purple) mite, citrus rust mite, six-spotted mite, Texas citrus mite, Yuma mite. Apply at the rate of 50 to 75 lbs. per acre. Application may be made up to the day of harvest. If additional applications are necessary, do not apply within 30 days of the previous application. When applied by aircraft, craft should fly over the top of each tree row to insure adequate coverage. When applied by ground machine, it should be applied to both sides of the tree to insure adequate coverage. Do not apply to kumquat or citron.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

50 LB. NET

DESTROY EMPTY CONTAINER — BURN — STAY OUT OF SMOKE

USDA Reg. No. 476-1899

Made in U.S.A.
Stauffer Chemical Company, Inc.

W WE 680521 SW