

OBSERVE THE FOLLOWING USE PRECAUTIONS

Highly toxic to fish and wildlife. Do not contaminate any body of water nor apply where runoff could occur.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues will not exceed established tolerances, use only at recommended rates.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.)

DIRECTIONS FOR USE

SPRAYING

Pour recommended amount of this material into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to animals may result. Do not combine with other materials.

LIVESTOCK OILERS OR BACKRUBBERS

For control of horn flies and as an aid to reduce lice infestations on beef cattle and swine, dilute this material at the rate of 1 gallon to 10 of diesel fuel. Place the mixture in oil reservoir or on burlap or oiler device and keep wet at all times. Place livestock oiler equipment where animals have free access to rubbing. Primarily near watering facilities or wherever animals congregate. Do not use on dairy animals or animals under 6 months of age. Do not permit animals access to treatment within 30 days of slaughter.

RESIDUAL SPRAY

For control of House Flies, Mosquitoes and Stable Flies. Use 1 gal. of this material to 9 gals. of water. Apply spray at the rate of 1/2 to 1 gal. per 1000 sq. ft. of surface. Thoroughly treat exteriors of barns, stock sheds and other farm buildings as well as manure, vegetation and refuse. Do not use in dairy barns or poultry houses. Repeat as needed.

LOT _____

ACCEPTED

8/12/63
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON, REGISTERED UNDER NO. 476-1913 SUBJECT TO ATTACHED COMMENTS.



**LINDANE TOXAPHENE
LIVESTOCK SPRAY AND
OILER CONCENTRATE .2-4E**

For Use on Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange and Ticks.

Contains .2 Lbs. Lindane and 4 Lbs. Toxaphene Per Gallon

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride (from Lindane)	2.2%
Toxaphene (Technical chlorinated camphene)*	45.1%
Petroleum hydrocarbon solvent	46.0%

INERT INGREDIENTS:..... 6.7%

* Chlorine Content 67% to 69% **100.0%**

WARNING Keep Out of Reach of Children
MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN.
Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not contaminate feed and food. In case of contact, wash immediately with soap and water. Do not use on household pets or humans.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME
Protect From Temperatures Below 0°F.

GAL. NET

STAUFFER CHEMICAL COMPANY

NEW YORK

RECOMMENDATIONS

Pest

Beef Cattle
Horn Flies, Lice,
Sarcoptic Mange, Tick

Swine
Lice, Sarcoptic Mange

Sheep
Lice, Keds (Sheep Tick),
Psoroptic or Sarcoptic
Scabs, Ticks

APPLICATION

Horn Flies—spray back
Lice, Ticks, Sarcoptic
mange thoroughly. Repeat

CAUTION

Do not treat sick animals
of any age. Do not spray at
slaughter. Do not treat
access to the shade fold
when weather is excessive
farrowing, nor for at least

NOTICE: Stauffer Chemical Company warrants the material, except the

USDA Reg. No. 476-1913

ACCEPTED
 8/12/03
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 177,195 SUBJECT
 TO ATTACHED COMMENTS.



LINDANE TOXAPHENE .2-4E

LIVESTOCK SPRAY AND OILER CONCENTRATE

For Use on Beef Cattle, Sheep and Swine. Controls Horn Flies, Lice, Sarcoptic and Psoroptic Mange and Ticks.

Contains .2 Lbs. Lindane and 4 Lbs. Toxaphene Per Gallon

ACTIVE INGREDIENTS:

Gamma isomer of benzene hexachloride (from Lindane)	2.2%
Toxaphene (Technical chlorinated camphene)*	45.1%
Petroleum hydrocarbon solvent	46.0%
INERT INGREDIENTS:	6.7%
* Chlorine Content 67% to 69%	100.0%

WARNING **Keep Out of Reach of Children**
MAY BE FATAL IF SWALLOWED. MAY BE ABSORBED THROUGH SKIN.
 Do not breathe spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not contaminate feed and food. In case of contact, wash immediately with soap and water. Do not use on household pets or humans.

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME
 Protec From Temperatures Below 0°F.

1 GAL. NET
STAUFFER CHEMICAL COMPANY
 NEW YORK

RECOMMENDATIONS

Pest	50 Gal. of Water	5 Gal. of Water
	Amount of Concentrate to Use Per:	
<u>Beef Cattle</u>		
Horn Flies, Lice,	2 qt. for Sprays	4/5 cup
Sarcoptic Mange, Ticks	2 qt. for Dips	4/5 cup
<u>Swine</u>		
Lice, Sarcoptic Mange	2 qt. for Sprays	4/5 cup
	2 qt. for Dips	4/5 cup
<u>Sheep</u>		
Lice, Keds (Sheep Tick), Psoroptic or Sarcoptic Sheep Scabs, Ticks	2 qt. for Sprays	4/5 cup
	1 qt. for Dips	2/5 cup

APPLICATION

Horn Flies—spray back of animals every 3 weeks.
 Lice, Ticks, Sarcoptic Mange and Sarcoptic Sheep Scab—Immerse or spray animals thoroughly. Repeat after 2-3 weeks if needed.

CAUTION

Do not treat sick animals. Do not use on dairy animals or animals under 6 months of age. Do not spray animals within 30 days of slaughter or dip within 60 days of slaughter. Do not treat emaciated or lactating animals. Animals should have access to the shade following dipping or spraying in warm weather. Do not treat when weather is excessively cold or hot. Do not treat sows within two weeks of farrowing, nor for at least 3 weeks thereafter.

NOTICE: Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-1913 Made in U.S.A. Stauffer Chemical Company, New York 3-SF-680516-NP