

476-1917

8-15-1972

1/2

IMIDAN 50-WP

EPA Reg. No. 476-1917

Supplemental Label Recommendations**ACCEPTED**

AUG 15 1972

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 476-1917 SUBJECT
 TO ATTACHED COMMENTS.

DECIDUOUS FRUIT -- FOLIAR SPRAYSApples:

For apple maggot, codling moth, elm spanworm, green apple aphid, fruit tree leafroller, gypsy moth, plum curculio, red banded leafroller, rosy apple aphid control and tarnished plant bug suppression, apply 1 to 1-1/2 pounds per 100 gallons water (or a maximum of 8 pounds per acre per application) as a full cover spray. Use 1-1/2 pounds per 100 gallons water for plum curculio. Repeat as necessary in accordance with insect infestations and local or state spray programs. Usually suppresses European red and two spotted mites when used in a seasonal program. Do not apply within 7 days of harvest. Recommended for control of elm spanworm, gypsy moth and fruit tree leafroller in the Northeast only.

Pears:

For codling moth, elm spanworm, fruit tree leafroller, green apple aphid, gypsy moth, pear psylla, plum curculio, red banded leafroller, rosy apple aphid control and tarnished plant bug suppression, apply 1 to 1-1/2 pounds per 100 gallons water (or a maximum of 10 pounds per acre per application) as a full cover spray. Use 1-1/2 pounds per 100 gallons water for plum curculio. Repeat as necessary in accordance with insect infestations and local or state spray programs. Usually suppresses European red and two spotted mites when used in a seasonal program. Do not apply within 7 days of harvest. Recommended for control of elm spanworm, fruit tree leafroller and gypsy moth in the Northeast only.

DECIDUOUS SHADE AND ORNAMENTAL TREES AND WOODY EVERGREENS

Note: Limit all uses for control of gypsy moth, elm spanworm, spring cankerworm and birch leaf miner to the Northeastern U. S. only.

IMIDAN 50-WP is recommended for use by commercial applicators on deciduous shade and ornamental trees and woody evergreens in parks, recreational areas, along thoroughfares and other localized areas where infestations of gypsy moth, spring cankerworm, elm spanworm or birch leaf miner threaten defoliation.

•••••
 •••••
 •••••

•••••
 •••••
 •••••

- . Recommended on Deciduous Shade and Ornamental Trees (ash, beech, oak, dogwood, willow, hickory, hawthorne, birch, elm, maple) to control: elm spanworm, spring cankerworm, gypsy moth.
- . Recommended on Woody Evergreens (arborvitae, cedar, fir, hemlock, juniper, pine, spruce, yew) to control: gypsy moth.
- . Recommended on Birch Trees to control: birch leaf miner.

When such insects or their damage occur, apply at a rate of 1-1/2 pounds IMIDAN 50-WP per 100 gallons of water, thoroughly wetting all parts of the affected plants to the point of runoff. Repeat application as necessary to maintain insect control.

Choose a cool, calm period, preferably in early morning or evening. Do not apply if rain is expected or before leaf surfaces are dry.

ARC 720621-0

