


# IMIDAN 50-WP

EPA Reg. No. 476-1917

## NEW LABEL RATE ON ALFALFA IN CALIFORNIA

**ALFALFA (California Only):** For control of Common Alfalfa Weevil Larvae and Egyptian Alfalfa Weevil Larvae, apply 1-1/2 pounds of IMIDAN 50-WP in 10 to 50 gallons of water per acre (20 to 50 gallons of water for dense stands) by ground equipment, or in 5 to 10 gallons of water per acre by aircraft. Consult your local Farm Advisor regarding the proper timing of application. Larvae should be sprayed when they are actively feeding. Do not apply more than once per cutting. Do not graze or cut for hay within 14 days of application. Do not use a spreader, sticker, or surfactant with IMIDAN 50-WP on alfalfa.

1-11-80



# of 4

Imidan Cases

**ALFALFA (California Only):** For control of Common Alfalfa Weevil Larvae and Egyptian Alfalfa Weevil Larvae, apply 1-1/2 pounds of IMIDAN 50-WP in 10 to 50 gallons of water per acre (20 to 50 gallons of water for dense stands) by ground equipment, or in 5 to 10 gallons of water per acre by aircraft. Consult your local Farm Advisor regarding the proper timing of application. Larvae should be sprayed when they are actively feeding. Do not apply more than once per cutting. Do not graze or cut for hay within 14 days of application. Do not use a spreader, sticker, or surfactant with IMIDAN 50-WP on alfalfa.

1-11-80

Imidan 30 lb. Bags

**ALFALFA (California Only):** For control of Common Alfalfa Weevil Larvae and Egyptian Weevil Larvae, apply 1-1/2 pounds of IMIDAN 50-WP in 10 to 50 gallons of water per acre (20 to 50 gallons of water for dense stands) by ground equipment, or in 5 to 10 gallons of water per acre by aircraft. Consult your local Farm Advisor regarding the proper timing of application. Larvae should be sprayed when they are actively feeding. Do not apply more than once per cutting. Do not graze or cut for hay within 14 days of application. Do not use a spreader, sticker, or surfactant with IMIDAN 50-WP on alfalfa.

1-11-80

Imidan 5 lb. Bags