


# CAPTAN-CYTHON-METHOXYCHLOR 65-1.5-10 SEED PROTECTANT

AGRICULTURAL FUNGICIDE-INSECTICIDE  
FOR SEED FUNGI DISEASE AND INSECT CONTROL



CAPTAN-CYTHON-METHOXYCHLOR 65-1.5-10 SEED PROTECTANT is a special wettable powder formulation for the slurry treatment of seeds, prior to planting, for protection against certain fungus diseases and insects. This material incorporates the use of Premium Grade Malathion.

## DIRECTIONS FOR USE

**SLURRY SEED TREATMENT:** Pre-mix the recommended amount of this material with a little water in a bucket and make a thin paste. Stir this paste into the required amount of water for the slurry equipment to be used.


**NOTE:** Seed treated by the slurry method should not be bagged until such time as the seed has had an opportunity to dry. Store in well ventilated areas. If seed is bagged when wet, heat is developed and impairs seed germination.

**IMPORTANT:** Treated seed must not be used for human consumption nor as animal feed. Bags containing treated seed should be labeled:

TREATED SEED

DO NOT USE FOR FEED, FOOD OR OIL PURPOSES

The Federal Government and some states have issued regulations for the application of special tags or labels to be affixed to bags containing treated seed.



### ACTIVE INGREDIENTS:

Captan*	65.0%
CYTHON**	1.5%
Methoxychlor, Technical***	10.0%
INERT INGREDIENTS:	23.5%
	100.0%

\*N-(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

### \*\* Malathion

O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate from Premium Grade Malathion.

\*\*\*Equivalent to 8.8% of 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.2% other isomers and related compounds.

NOT LICENSED FOR USE IN COUNTRIES WHERE VALID PATENTS STILL EXIST.

## CAUTION

Keep Out of Reach of Children

Harmful if swallowed, inhaled or absorbed through skin.

Avoid prolonged or repeated contact with skin.

Wash with soap and water after handling.

Do not contaminate feed or food.

Toxic to fish. Do not contaminate any body of water nor apply where runoff could occur.

Do not contaminate water by cleaning of equipment or disposal of wastes.

PROTECT FROM EXCESSIVE HEAT • STORE IN A COOL DRY PLACE  
CONTAINER SHOULD BE DISPOSED BY CRUSHING AND BURYING WITH WASTES

## RECOMMENDATIONS

For the control of seed rot, damping-off diseases, and kernel smut of sorghum, and for the protection of seed against Cadelle, Confused Flour Beetle, Flat Grain Beetle, Granary Weevil, Lesser Grain Borer, Red Flour Beetle, Rice Weevil, Rusty Grain Beetle, and Saw-toothed Beetle.

AMOUNT OF CAPTAN-CYTHON-METHOXYCHLOR  
65-1.5-10 TO USE ON SEEDS

SEED	SLURRY TREATMENT
Corn (field) seed	1 2/3 oz./cwt. or 1 oz./bu.
Sorghum (hull-less) seed	3 1/2 oz./cwt. or 1 3/4 oz./bu.

Further specific information on this subject may be obtained from State Agricultural Experiment Stations or the State Agricultural Extension Service.

**NOTICE:** Stauffer Chemical Company makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on Stauffer's label.

USDA Reg. No. 476-1918 Made in U.S.A. OOM680612MW

50 LB. NET