# DUSTING SULFUR 50%

## **AN AGRICULTURAL INSECTICIDE and FUNGICIDE**

| ACTIVE INGREDIENT |            |       |     |
|-------------------|------------|-------|-----|
| Sulfur            | <b>.</b> . | · .   | 50% |
| INERT INGREDIENTS | <b>.</b> . | ••••• | 50% |
|                   |            |       |     |

## CAUTION

100% Ap

Sulfur will cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on apricots. Anjou and comice pears, cranberries, cucurbits (cucumbers, squash and some varieties of melons), filberts, spinach, tung trees walnuts and other sensitive plants.

Except on cotton, do not apply when shade temperatures exceed or are likely to exceed 90 F. Sulfur may burn foliage and fruit during periods of high temperatures and under certain climatic conditions.

Do not use sulfur within 30 days of an oil spray (60 days on citrus).

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

NOTICE: Stauffer Chemical Company makes no warranty of merchantability or any other express or implied warranty concerning this material. It shall not be held responsible for personal injury, property damage, or other loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability resulting from such handling, storage or use.

BUR FERETY BAG - STAN JUT OF SHOKE

NET WEIGHT 50 POUNDS

## RECOMMENDATIONS

#### FRUIT CROPS

Apples: Scab, powdery mildew — apply 60 lbs. per acre in the green tip to pink bud stage. Use on sulfur-tolerant varieties only.

Strawberries: Powdery mildew - apply 40 lbs. per acre.

#### **VEGETABLE AND FIELD CROPS**

Alfalfa: Spider mites — apply 40 lbs. per acre. Caution. sulfur may cause petal burn to open flowers

**Beans:** Powdery mildew, rust, spider mites — apply 45 lbs. per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. Caution: may cause injury to some varieties of beans.

Clover: Powdery mildew, spider mites --- apply 40 lbs. per acre.

Peas: Powdery mildew --- apply 40 lbs. per acre.

Sugar beets: Spider mite - apply 50 lbs. per acre.

Grass Seed Crops: Timothy mite — apply 50 lbs. per acre

